

RADIATION FIELD FOR EINSTEIN VACUUM EQUATIONS WITH SPACIAL DIMENSION $n \geq 4$

FANG WANG

1. INTRODUCTION

The Einstein vacuum equations determine a manifold M^{1+n} with a Lorentzian metric with vanishing Ricci curvature:

$$(1.1) \quad R_{\mu\nu} = 0.$$

The set $(\mathbb{R}_{t,x}^{1+n}, m)$: standard Minkowski $m = -dt^2 + \sum_{i=1}^n dx^i dx^i$ on $\mathbb{R}_{t,x}^{1+n}$ describes the Minkowski space-time solution of the system (1.1). The problem of stability of Minkowski space appears in Cauchy formulation of the Einstein Vacuum equations: for a given n dimensional manifold Σ^n with a Riemannian metric g_0 and a symmetric two tensor k_0 , such that $(\Sigma^n; g_0, k_0)$ is close to $(\mathbb{R}^n; m_e, 0)$ where m_e the Euclidean metric, we want to find a $n+1$ dimensional manifold M^{1+n} with a Lorentzian metric g of signature $(n, 1)$ satisfying the Einstein vacuum equations (1.1), and an embedding $\Sigma^n \subset M^{1+n}$ such that g_0 is the restriction of g to Σ and k_0 is the second fundamental form of (Σ^n, g_0) in (M^{1+n}, g) . The Cauchy problem is overdetermined which imposes compatibility condition on the Cauchy data: the constraint equations

$$(1.2) \quad R_0 - [k_0]_j^i [k_0]_i^j + [k_0]_i^i [k_0]_j^j = 0, \quad \nabla^j [k_0]_{ij} - \nabla_i [k_0]_j^j = 0, \forall i = 1, \dots, n.$$

Here R_0 is the scalar curvature of g_0 and ∇ is covariant differentiation w.r.t. g_0 .

The seminal result of Choquet-Bruhat [CB1] followed by the work [CBG] showed existence and uniqueness up to diffeomorphism of a maximal globally hyperbolic smooth space-time arising from any set of smooth initial data. The global stability of Minkowski space for the Einstein vacuum equations was shown in a remarkable work of Christodoulou-Klainerman for strongly asymptotic flat initial data [CK]. The approach taken in that work viewed the Einstein vacuum equations as a system of equations

$$D^\alpha W_{\alpha\beta\gamma\delta} = 0, \quad D^\alpha * W_{\alpha\beta\gamma\delta} = 0$$

for the Weyl tensor $W_{\alpha\beta\gamma\delta}$ of the metric $g_{\alpha\beta}$ and used generalized energy inequalities associated with the Bel-Robinson energy-momentum tensor, constructed from components of W , and special geometrically constructed vector fields, designed to mimic the rotation and the conformal Morawetz vector fields of the Minkowski space-time, i.e. "almost conformal Killing" vector fields of the unknown metric g . The proof was manifestly invariant.

The Einstein vacuum equations are invariant under diffeomorphism. In the work of Choquet-Bruhat, this allows her to choose a special *harmonic gauge*, also referred as *wave coordinates*, in which the Einstein vacuum equations become a system of quasilinear wave equations on the components of the unknown metric $g_{\mu\nu} = m_{\mu\nu} + h_{\mu\nu}$:

$$(1.3) \quad \tilde{\square}_g h_{\mu\nu} = F_{\mu\nu}(h)(\partial h, \partial h),$$

where the reduced wave operator is defined by $\tilde{\square}_g = g^{\alpha\beta} \partial_\alpha \partial_\beta$ and $F(u)(v, v)$ depends quadratically on v and analytically on u for u small. Wave coordinates $\{x^\mu\}$ are required to be solutions of the wave equations

$$\square_g x^\mu = 0, \quad \text{for } \mu = 0, 1, \dots, n,$$

where the geometric wave operator is defined by $\square_g = D_\alpha D^\alpha = g^{\alpha\beta} \partial_\alpha \partial_\beta - g^{\alpha\beta} \Gamma_{\alpha\beta}^\nu \partial_\nu$. The metric $g_{\mu\nu}$ relative to wave coordinates $\{x^\mu\}$ satisfies the *harmonic gauge condition*:

$$(1.4) \quad \Gamma_\mu = 0, \quad \text{for } \mu = 0, 1, \dots, n,$$

where $\Gamma_\mu = g_{\mu\nu}\Gamma^\nu = g_{\mu\nu}g^{\alpha\beta}\Gamma_{\alpha\beta}^\nu = g^{\alpha\beta}\partial_\beta g_{\alpha\mu} - \frac{1}{2}g^{\alpha\beta}\partial_\mu g_{\alpha\beta}$. Under this condition the geometric wave operator $\square_g$ is equal to the reduced wave operator $\tilde{\square}_g$ and the stability of Minkowski space can reformulate as follows: given a pair of small symmetric two tensors (h^0, h^1) on $\{t=0\} \subset \mathbb{R}_{t,x}^{1+n}$ which satisfies the harmonic gauge conditions:

$$(1.5) \quad \Gamma_\mu|_{t=0} = 0, \quad \partial_t \Gamma_\mu|_{t=0} = 0,$$

we want to find a Lorentian metric $g = m + h$ of signature $(n, 1)$ satisfying the reduced Einstein equations (1.3) such that $(h|_{t=0}, \partial_t h|_{t=0}) = (h^0, h^1)$.

Theorem 1 (H. Lindblad, I. Rodnianski). *Given Cauchy data (h^0, h^1) in coordinates (t, x) satisfying the harmonic gauge conditions (1.5), then the solution to the reduced Einstein equations (1.3) provides a solution to the Einstein vacuum equations (1.1).*

The use of harmonic gauge goes back to the work of Einstein on post-Newtonian and post-Minkowski expansions. In the PDE terminology the result of Choquet-Bruhat corresponds to the *local well-posedness* of the Cauchy problem for the Einstein vacuum equations with smooth initial data. The reduced Einstein vacuum equations satisfies the *null condition* for spacial dimension $n \geq 4$, which ensures the global existence theorem for small Cauchy data. The concept of null condition was designed to detect systems for which solutions are asymptotically free and decay like solutions of a linear equation. See [Ch1], [Kl] and [Ho1]. For $n = 3$, it can be shown that the Einstein vacuum equations in harmonic gauge do not satisfy the null condition. Moreover, Choquet-Bruhat [CB2] showed that even without imposing a specific gauge the Einstein equations violate the null condition. However, the Einstein vacuum equations in harmonic gauge satisfy the *weak null condition* and recently Lindblad-Rodnianski reproved the global existence for Einstein vacuum equations in harmonic gauge for general asymptotic flat initial data in harmonic gauge by combining it with the vector field method [LR1] [LR2].

In this paper, we apply the radiation field theory due to Friedlander to study the asymptotic behavior of solutions to Einstein vacuum equations in harmonic gauge. Friedlander's radiation field was used by Hörmander to study the asymptotic behavior of solutions to linear hyperbolic equation in the following coordinates:

$$\rho = \frac{1}{|x|}, \quad \tau = t - r, \quad \theta = \frac{x}{r},$$

for $|x|$ large. For instance, consider the Cauchy problem in Minkowski space-time as follows:

$$\square_m u(t, x) = 0, \quad u(0, x) = u_0(x), \quad \partial_t u(0, x) = u_1(x),$$

where $u_0, u_1 \in C_c^\infty(\mathbb{R}^n)$. Near $\rho = 0$, by writing $u = \rho^{\frac{n-1}{2}} \tilde{u}$ and studying the equivalent equation

$$(\square_{\tilde{m}} + \frac{(n-1)(n-3)}{4})\tilde{u} = 0$$

with $\tilde{m} = \rho^2 m$, Friedlander showed that $\tilde{u}$ is smooth up to $\rho = 0$. Then the radiation field is the image of the map

$$\mathcal{R}_{\mathcal{LP}} : \dot{H}^1(\mathbb{R}^n) \times L^2(\mathbb{R}^n) \ni (u_0, u_1) \longrightarrow \partial_\tau \tilde{u}|_{\rho=0} \in L^2(\mathbb{R}_\tau \times S_\theta^{n-1}),$$


which is an isometric isomorphism. Here $\mathcal{R}_{\mathcal{LP}}$ is essentially the *Møller wave operator* and also the free space translation representation of Lax and Phillips.

I generalize this idea by considering the conformal transformation of the reduced Einstein vacuum equations on a suitable compactification of $\mathbb{R}_{t,x}^{1+n}$ as follows. Here $S_1^\pm$ is the compactification of null infinity and $\Sigma_0 = \{t=0\}$ is Cauchy surface. Let $\tilde{\rho}$ be the total boundary defining function and ρ_0, ρ_1, ρ_2 be the defining functions for corresponding boundary hypersurfaces $S_0, S_1^\pm, S_2^\pm$. With a conformal change $\tilde{g} = \tilde{\rho}^2 g$ and $\tilde{h} = \tilde{\rho}^{\frac{1-n}{2}} h$, the reduced Einstein equations (1.3) are equivalent to

$$(1.6) \quad (\square_{\tilde{g}} + \gamma(\tilde{h}))\tilde{h}_{\mu\nu} = \rho_1^{\frac{n-5}{2}} (\rho_0 \rho_2)^{\frac{n-1}{2}} \tilde{F}_{\mu\nu}(\tilde{h}, \tilde{h}),$$

where $\gamma(\tilde{h}) = -\tilde{\rho}^{\frac{n-1}{2}} \square_{\tilde{g}} \tilde{\rho}^{\frac{1-n}{2}}$ and $\tilde{F}_{\mu\nu}(\tilde{h}, \tilde{h}) = \rho_1^{1-n} (\rho_0 \rho_2)^{-1-n} F_{\mu\nu}(\tilde{\rho}^{\frac{n-1}{2}} \tilde{h}, \tilde{\rho}^{\frac{n-1}{2}} \tilde{h})$. See Section 5 for more details. Denote by $\tilde{\mathcal{V}}_\epsilon^{N,\delta}$ the space of elements in the weighted b-Sobolev space

$$\rho_0^{\frac{n-1}{2}+\delta} H_b^{N+1}(\Sigma_0) \times \rho_0^{\frac{n+1}{2}+\delta} H_b^{N+1}(\Sigma_0)$$

FIGURE 1. the blown-up space $X = [\overline{\mathbb{R}_{t,x}^{1+n}} : \overline{\{|t| = |x|\}} \cap \overline{\partial \mathbb{R}_{t,x}^{1+n}}]$

with norm less than ϵ and satisfying the harmonic gauge conditions (1.5), for some $N \geq n + 6$, $\frac{1}{2} > \delta > 0$ and $\epsilon > 0$ small. Then by energy estimate method I show that for $n \geq 4$, if $(h^0, h^1) \in \tilde{\mathcal{V}}_\epsilon^{N,\delta}$ then $\tilde{h} = (\rho_0 \rho_1 \rho_2)^{\frac{1-n}{2}} h$ is $C^{0,\delta}$ up to $S_1^\pm$ and hence the radiation field for h is well defined which satisfies the corresponding harmonic gauge conditions:

$$(1.7) \quad \tilde{\Gamma}_\mu|_{S_1^\pm} = 0, \quad \text{for } \mu = 0, 1, \dots, n,$$

where $\tilde{\Gamma}_\mu = \tilde{\rho}^{\frac{1-n}{2}} \Gamma_\mu$. Combining this with the linear radiation field theory and the implicit function theorem, we further show that $\tilde{h}|_{S_1^\pm}$ also lies in some weighted b-Sobolev space on $S_1^\pm$. More explicitly, denote by $\tilde{\mathcal{W}}_\epsilon^{N,\delta}$ the elements in

$$(\rho_0 \rho_2)^\delta [H_b^{\frac{1}{2}-\delta}(\mathbb{R}; H^{N+\frac{1}{2}+\delta}(\mathbb{S}^{n-1})) \cap L^2(\mathbb{S}^{n-1}; H_b^{N+1}(\mathbb{R}))]$$

where we take the equivalence $S_1^\pm = \overline{\mathbb{R}_\tau} \times \mathbb{S}_\theta^{n-1}$, with norm less than ϵ' and satisfying (1.7). Then $\tilde{h}|_{S_1^\pm} \in \tilde{\mathcal{W}}_\epsilon^{N,\delta}$ for some $\epsilon' > 0$. Here both $\tilde{\mathcal{V}}_\epsilon^{N,\delta}$ and $\tilde{\mathcal{W}}_\epsilon^{N,\delta}$ are Banach manifolds. See Section 2 for the definition of weighted b-Sobolev spaces and Section 3, 7 for more details about $\tilde{\mathcal{V}}_\epsilon^{N,\delta}$ and $\tilde{\mathcal{W}}_\epsilon^{N,\delta}$.

The main theorem of this paper is the following:

Theorem 2. *For $n \geq 4$, $N \geq n + 6$, $\frac{1}{2} > \delta > 0$ and $\epsilon > 0$ small, if $(h^0, h^1) \in \tilde{\mathcal{V}}_\epsilon^{N,\delta}$, then the Einstein equation (1.1) has a global solution h such that $\tilde{h}$ is $C^{0,\delta}$ up to $S_1^\pm$. Hence the Radiation field is well defined and the nonlinear Møller wave operator defines a continuous and open map*

$$\mathcal{R}_\epsilon^\pm : \tilde{\mathcal{V}}_\epsilon^{N,\delta} \ni (h^0, h^1) \longrightarrow \tilde{h}|_{S_1^\pm} \in \tilde{\mathcal{W}}_{C\epsilon}^{N,\delta}$$

for some constant $C > 0$.

The open property guarantees the solution for characteristic initial value problem for Einstein Vacuum equations.

Corollary 1. *Given characteristic data $\tilde{h}^{S_1^+} \in \tilde{\mathcal{W}}_\epsilon^{N,\delta}$ for some $N \geq n + 6$, $\delta \in (0, \frac{1}{2})$ and $\epsilon > 0$ small enough, there exists an asymptotically flat solution g to the Einstein vacuum equation (1.1) such that $\tilde{\rho}^{\frac{1-n}{2}}(g - m)|_{S_1^+} = \tilde{h}^{S_1^+}$. Moreover, (t, x) are wave coordinates w.r.t. g globally.*

For $n = 3$, the Cauchy data of interest for reduced Einstein vacuum equations has an asymptotic leading term $-M|x|^{-1}\delta_{ij}$ for some constant $M > 0$ small, which has a long range effect at null infinity. Since $-M|x|^{-1}\delta_{ij}$ provides a solution to the linearization of (1.3) in a neighborhood of null infinity in X , we may expect that the essential change of the geometry of perturbed solution $g = m + h$ only comes from the asymptotic leading term, i.e. the constant M . By a different change of coordinates to $\tau = t - |x| - M \log |x|$, which was suggested by Friedlander to study the linear equation $\square_g u = 0$ with such background metric, I find a corresponding compactification of $\mathbb{R}_{t,x}^{1+n}$ such that the radiation field is well defined in a similar way. This will be discussed in details in a separate paper.

The outline the the paper is as follows: in Section 2 we choose a compactification of $\mathbb{R}_{t,x}^{1+n}$ and give basic definitions used through out the paper; in Section 3 and Section 4, we study the solution space of constraint equations (1.2) and of harmonic gauge conditions (1.5) separately; in Section 5, we make a conformal transformation of the reduced Einstein equations and study the metric and wave operator in the conformal setting; in Section 6, we do energy estimates for the equations (1.6) and show the existence of radiation field; in Section 7, we modify the result obtained in Section 6 by applying the linear theory and implicit function theorem and obtain the isomorphism property for the nonlinear Møller wave operator.

Acknowledgment: I want to thank to my adviser Richard Melrose for inspirations and valuable discussion on this topic. Without his help, this paper would not exist. I also would like apologize to all who expressed interest in this work for the long delay between the first announcement of these results and the appearance of this monograph.

2. GEOMETRIC SETTING

This section is a preparation for describing the problem and dealing with it in a manifold with corners arising from a suitable compactification of $\mathbb{R}_{t,x}^{1+n}$. To simplify the notation, we take $t = x^0$, $x = (x^1, \dots, x^n)$ and use the lower case English alphabet $i, j, k, l, \dots$ as indices taking value in $\{1, 2, \dots, n\}$ and the Greek alphabet $\alpha, \beta, \mu, \nu, \dots$ as indices taking value in $\{0, 1, 2, \dots, n\}$. Moreover, we use the capital English alphabet $I, J, K, L, \dots$ as multi indices. The change between superscript and subscript is taken over Minkowski metric $m = -dt^2 + \sum_{i=1}^n dx^i dx^i$. For example, $x_0 = -x^0$, $x_i = x^i$.

2.1. Compactification of the Space-time. The usual radial compactification of $\mathbb{R}_{t,x}^{1+n}$, denoted by X_0 , can be realized by embedding

$$\Phi : \mathbb{R}_{t,x}^{1+n} \ni (t, x) \longrightarrow \left(\frac{1}{\sqrt{1+t^2+|x|^2}}, \frac{t}{\sqrt{1+t^2+|x|^2}}, \frac{x}{\sqrt{1+t^2+|x|^2}} \right) \in \mathbb{R}^{2+n}$$

and taking $X_0 = \overline{\Phi(\mathbb{R}_{t,x}^{1+n})}$, which is a closed half sphere of dimension $n+1$. Then all null rays w.r.t. Minkowski metric converge to an embedded submanifold of X_0 on the boundary:

$$L = \overline{\Phi(\{|t| = |x|\})} \cap \partial X_0 = \{(0, \pm \frac{1}{\sqrt{2}}, \pm \frac{x}{\sqrt{2}}) : x \in \mathbb{R}^n, |x| = 1\}.$$

The compactification space X , which we use in this paper to study the asymptotic behavior of Einstein vacuum solutions that are close to Minkowski metric, is X_0 blown up L . Hence X is a manifold with corners up to codimension 2. See Figure 1. More explicitly, let $d(p, q)$ be the distance function on X_0 w.r.t. standard spherical metric and $U_\epsilon = \{p \in X_0 : d(p, L) < \epsilon\}$; let

$$\Psi : X_0 \setminus L \longrightarrow X_0 \setminus U_{\frac{1}{4}}$$

defined as follows: if $d(p, L) > \frac{1}{2}$ then $\Psi(p) = p$; if $d(p, L) \leq \frac{1}{2}$ then

$$d(\Psi(p), L) = d(\Psi(p), p) + d(p, L), \quad d(\Psi(p), L) = \psi(d(p, L))$$

where $\psi \in C^\infty([0, \frac{1}{2}])$ satisfying $\psi(0) = \frac{1}{4}$, $\psi(\frac{1}{2}) = \frac{1}{2}$, $\psi' > 0$ and $\psi'(d) = 1$ for $d \in [\frac{3}{8}, \frac{1}{2}]$. Then the blowup can be realized by taking

$$(2.1) \quad X = \overline{\Psi(X_0 \setminus L)} = X_0 \setminus U_{\frac{1}{4}} \subset \mathbb{R}^{2+n}.$$

Obviously, $\Psi\Phi$ embeds $\mathbb{R}_{t,x}^{1+n}$ into $\mathbb{R}^{2+n}$ with image $\mathring{X}$, the interior of X . This is closely related to Penrose's diagram. The three essentially different boundary hypersurfaces can be represented with the three regions at infinity: spatial infinity S_0 , null infinity $S_1^\pm$ and temporal infinity $S_2^\pm$ where the latter have forward and backward components. In particular, we can identify $S_0, S_1^\pm$ with $\mathbb{R} \times \mathbb{S}^{n-1}$, where $\mathbb{R}$ is the radial compactification of $\mathbb{R}$, and $S_2^\pm$ with closed ball of dimension n . Denote

$$M_1(X) = \{S_0, S_1^\pm, S_2^\pm\}.$$

Definition 1. For $S \subset M_1(X)$, we say ρ_S a defining function for S iff

$$0 \leq \rho_S \in C^\infty(X), \quad S = \{\rho_S = 0\}, \quad d\rho_S \neq 0.$$

Two boundary defining functions ρ_S, ρ'_S are equivalent on a domain $\Omega \subset X$ iff

$$\rho_S = e^\psi \rho'_S \text{ for some } \psi \in C^\infty(\bar{\Omega}).$$

We say $\tilde{\rho}$ a total boundary defining function for X iff

$$\tilde{\rho} = \prod_{S \in M_1(X)} \rho_S.$$

Throughout this paper, we use ρ_0, ρ_1, ρ_2 to denote the defining functions for $S_0, S_1^\pm, S_2^\pm$ and $\tilde{\rho} = \rho_0 \rho_1 \rho_2$ the total boundary defining function for ∂X . We make a special choice of them in local coordinates in the following.




FIGURE 2. the local coordinate charts on X

Choose a covering $\{\Omega_i : 0 \leq i \leq 5\}$ for X such that Ω_0 is a bounded domain in the interior $\overset{\circ}{X}$ and $\cup_{i=1}^5 \Omega_i$ covers ∂X . The corresponding local coordinates can be chosen from the following functions:

$$\begin{aligned} r &= |x|, \quad \tau = |t| - |x|, \quad \rho = \frac{1}{|x|}, \quad s = \frac{t}{|x|}, \quad \phi = \frac{1}{|t|}, \quad y = \frac{x}{|t|}, \\ a &= 1 - \frac{|t|}{|x|}, \quad b = \frac{1}{|x| - |t|}, \quad \bar{a} = 1 - \frac{|x|}{|t|}, \quad \bar{b} = \frac{1}{|t| - |x|}, \quad \theta^i = \frac{x^i}{|x|}, \quad \theta^0 = 1. \end{aligned}$$

More explicitly,

- In the domain $\Omega_0 = \{(t, x) : |t|^2 + |x|^2 < 1000\}$, choose

$$\tilde{\rho} = \rho_0 = \rho_1 = \rho_2 = 1.$$

- In the domain $\Omega_1 = \{(s, \rho, \theta) : -\frac{7}{8} < s < \frac{7}{8}, 0 \leq \rho < 1, \theta \in \mathbb{S}^{n-1}\}$, choose

$$\tilde{\rho} = \rho_0 = \rho, \quad \rho_2 = \rho_1 = 1.$$

- In the domain $\Omega_2 = \{(a, b, \theta) : 0 \leq a < \frac{7}{8}, 0 \leq b < 1, \theta \in \mathbb{S}^{n-1}\}$, choose

$$\tilde{\rho} = ab, \quad \rho_0 = b, \quad \rho_1 = a, \quad \rho_2 = 1.$$

- In the domain $\Omega_3 = \{(\tau, \rho, \theta) : -\tau_0 \leq \tau \leq \tau_0, 0 \leq \rho < 1, \theta \in \mathbb{S}^{n-1}\}$ for some $\tau_0 > 8$, choose

$$\tilde{\rho} = \rho_1 = \rho, \quad \rho_0 = \rho_2 = 1.$$

- In the domain $\Omega_4 = \{(\bar{a}, \bar{b}, \theta) : 0 \leq \bar{a} < \frac{7}{8}, 0 \leq \bar{b} < 1, \theta \in \mathbb{S}^{n-1}\}$, choose

$$\tilde{\rho} = \bar{a}\bar{b}, \quad \rho_1 = \bar{a}, \quad \rho_2 = \bar{b}, \quad \rho_0 = 1.$$

- In the domain $\Omega_5 = \{(\phi, y) : 0 \leq \phi < 1, |y| < \frac{7}{8}\}$, choose

$$\tilde{\rho} = \rho_2 = \phi, \quad \rho_1 = \rho_0 = 1.$$

Notice that, in the intersection of any two domains, the different choices for defining functions are equivalent. In many cases, for example when doing energy estimates, equivalent defining functions give equivalent statements. So we change the choice of ρ_i freely from one set of coordinates to the other in the rest of this paper if they give equivalent results. However, when performing energy estimates, we may not be able to restrict the discussion in a single coordinate chart and hence have to consider some of them together. See Section 6 for more details.

We also denote by $[X]^2$ a double space of X across the null infinity $S_1^+ \cup S_1^-$ such that $[X]^2$ is a manifold with boundary and the smooth structure on X extends to $[X]^2$, i.e. $C^\infty(X) = C^\infty([X]^2)|_X$. Here $[X]^2$ is not uniquely determined. Refer to [Me] for more details. The boundary of $[X]^2$ consists of three components, $[S_0]^2$, $[S_2^+]^2$ and $[S_2^-]^2$, which are also double spaces of S_0 , S_2^+ and S_2^- across their boundaries and hence are close manifolds. Moreover, $[S_0]^2$ can be identified with $\mathbb{S}^1 \times \mathbb{S}^{n-1}$ and $[S_2^\pm]^2$ can be identified with $\mathbb{S}^n$. We introduce $[X]^2$ for the reason that the conformal Minkowski metric, or with small perturbation, extends to a b-type Lorentz metric on $[X]^2$. See Section 4 for more details.

2.2. Vector Fields. Let $\mathcal{V}_b(X)$ (resp. $\mathcal{V}_b([X]^2)$) be the space of smooth vector fields on X which are tangent to ∂X (resp. $\partial[X]^2$). The relation between $\mathcal{V}_b([X]^2)|_X$ and $\mathcal{V}_b(X)$ is

$$\mathcal{V}_b([X]^2)|_X = \mathcal{V}_b(X) + C^\infty(X)V_1.$$

where $V_1 \in \mathcal{V}_b([X]^2)|_X$ transverse to $S_1^\pm$, i.e. for any $p \in S_1^\pm$, $V_1(p) \notin T_p S_1^\pm$. In particular, in domain $\Omega_2, \Omega_3, \Omega_4$, we can choose $V_1 = \partial_{\rho_1}$. Set

$$\begin{aligned} \partial_0 &= \partial_t, & \partial_i &= \partial_{x^i}, & \tilde{\partial}_i &= r(\delta_i^j - \theta_i \theta^j) \partial_j, \\ Z_{00} &= r \partial_r + t \partial_t, & Z_{0i} &= t \partial_i + x_i \partial_t, & Z_{ij} &= x_i \partial_j - x_j \partial_i. \end{aligned}$$

Here $\tilde{\partial}_i$ are the projection of ∂_i on $T\mathbb{S}^{n-1} \subset T\mathbb{R}^n$ satisfying $\theta^i \tilde{\partial}_i = 0$. We denote by $\tilde{Z}$ any vector field in $\{\partial_\mu, Z_{\mu\nu} : \mu, \nu = 0, \dots, n\}$ and $\tilde{\partial}^k$ any element in $\mathcal{V}_b^k(X)$ for $k \geq 1$.

Lemma 2.1. *The vector fields defined above satisfy the following properties:*

- (i) $\mathcal{V}(\mathbb{S}^{n-1}) = \text{Span}_{C^\infty(\mathbb{S}^{n-1})}\{Z_{ij} : i, j = 1, \dots, n\}$;
- (ii) $\mathcal{V}_b(X) = \text{Span}_{C^\infty(X)}\{\partial_\mu, Z_{\mu\nu} : \mu, \nu = 0, 1, \dots, n\}$;
- (iii) $\partial_\mu \in \rho_0 \rho_2 \mathcal{V}_b(X)$, $\mu = 0, \dots, n$.

Proof. The first property follows directly by $Z_{ij} = \theta_i \tilde{\partial}_j - \theta_j \tilde{\partial}_i \in T\mathbb{S}^{n-1}$ and $\tilde{\partial}_j = Z_{ij} \theta^i$. To prove (ii) and (iii), we only need to check them in local coordinates near ∂X :

- In Ω_1 with coordinates (s, ρ, θ) , $\mathcal{V}_b(\Omega_1) = \text{Span}_{C^\infty(\Omega_1)}\{\partial_s, \rho \partial_\rho, \tilde{\partial}_i : 1 \leq i \leq n\}$. Hence (ii) and (iii) follow by

$$\begin{aligned} \partial_t &= \rho \partial_s, \quad \partial_i = \rho(-\theta_i(\rho \partial_\rho + s \partial_s) + \tilde{\partial}_i), \\ Z_{00} &= -\rho \partial_\rho, \quad Z_{0i} = \theta_i(1 - s^2) \partial_s - \theta_i s \rho \partial_\rho + s \tilde{\partial}_i, \\ \implies \quad \partial_s &= (1 - s^2)^{-1}(Z_{0i} \theta^i - s Z_{00}), \quad \rho \partial_\rho = -Z_{00}. \end{aligned}$$

- In Ω_2 with coordinates (a, b, θ) , $\mathcal{V}_b(\Omega_2) = \text{Span}_{C^\infty(\Omega_2)}\{a \partial_a, b \partial_b, \tilde{\partial}_i : 1 \leq i \leq n\}$. Hence (ii) and (iii) follow by

$$\begin{aligned} \partial_t &= b(b \partial_b - a \partial_a), \quad \partial_i = -b \theta_i(b \partial_b - a \partial_a) + ab(-\theta_i a \partial_a + \tilde{\partial}_i), \\ Z_{00} &= -b \partial_b, \quad Z_{0i} = -\theta_i((2 - a)a \partial_a - b \partial_b) + (1 - a)\tilde{\partial}_i, \\ \implies \quad b \partial_b &= -Z_{00}, \quad a \partial_a = -(2 - a)^{-1}(Z_{00} + Z_{0i} \theta^i). \end{aligned}$$

- In Ω_3 with coordinates (τ, ρ, θ) , $\mathcal{V}_b(\Omega_3) = \text{Span}_{C^\infty(\Omega_3)}\{\partial_\tau, \rho\partial_\rho, \bar{\theta}_i : 1 \leq i \leq n\}$. Hence (ii) and (iii) follow by

$$\begin{aligned} \partial_t &= \partial_\tau, \quad \partial_i = -\theta_i \partial_\tau + \rho(-\theta_i \rho \partial_\rho + \bar{\theta}_i), \\ Z_{00} &= \tau \partial_\tau - \rho \partial_\rho, \quad Z_{0i} = -\theta_i(\tau \partial_\tau + (1 + \rho\tau)\rho \partial_\rho) + (1 + \rho\tau)\bar{\theta}_i, \\ \implies \quad \partial_\tau &= \partial_t, \quad \rho \partial_\rho = -(2 + \rho\tau)^{-1}(Z_{00} + Z_{0i}\theta^i). \end{aligned}$$

- In Ω_4 with coordinates $(\bar{a}, \bar{b}, \theta)$, $\mathcal{V}_b(\Omega_4) = \text{Span}_{C^\infty(\Omega_4)}\{\bar{a}\partial_{\bar{a}}, \bar{b}\partial_{\bar{b}}, \bar{\theta}_i : 1 \leq i \leq n\}$. Hence (ii) and (iii) follow by

$$\begin{aligned} \partial_t &= -\bar{b}(\bar{b}\partial_{\bar{b}} - \bar{a}\partial_{\bar{a}}) - \bar{a}\bar{b}(\bar{a}\partial_{\bar{a}}), \quad \partial_i = \bar{b}\theta_i(\bar{b}\partial_{\bar{b}} - \bar{a}\partial_{\bar{a}}) + \bar{a}\bar{b}(1 - \bar{a})^{-1}\bar{\theta}_i, \\ Z_{00} &= -\bar{b}\partial_{\bar{b}}, \quad Z_{0i} = -\theta_i((2 - \bar{a})\bar{a}\partial_{\bar{a}} - \bar{b}\partial_{\bar{b}}) + (1 - \bar{a})^{-1}\bar{\theta}_i, \\ \implies \quad \bar{b}\partial_{\bar{b}} &= -Z_{00}, \quad \bar{a}\partial_{\bar{a}} = -(2 - \bar{a})^{-1}(Z_{00} + Z_{0i}\theta^i). \end{aligned}$$

- In Ω_5 with coordinates (ϕ, y) , $\mathcal{V}_b(\Omega_5) = \text{Span}_{C^\infty(\Omega_5)}\{\phi\partial_\phi, \partial_{y^i} : 1 \leq i \leq n\}$. Hence (ii) and (iii) follow by

$$\begin{aligned} \partial_t &= -\phi(\phi\partial_\phi + y^i\partial_{y^i}), \quad \partial_i = \phi\partial_{y^i}, \\ Z_{00} &= -\phi\partial_\phi, \quad Z_{0i} = \partial_{y^i} - y_i(\phi\partial_\phi + y^j\partial_{y^j}), \\ \implies \quad \phi\partial_\phi &= -Z_{00}, \quad \partial_{y^i} = Z_{0i} - (1 - |y|^2)^{-1}y_i(Z_{00} - Z_{0j}y^j). \end{aligned}$$

□

2.3. Sobolev Spaces and Symbol Spaces. We define the b-type Sobolev space and conormal symbol space on X in the following. These spaces are used to characterize the regularity and asymptotic properties near boundary for Einstein vacuum solutions. Let m_0 be the Riemannian metric on X induced from Euclidean metric on $\mathbb{R}^{2+n}$ via (2.1) and denote by $dvol_{m_0}$ the corresponding volume form.

Definition 2. Define the b-Sobolev space $H_b^N(X)$ for any $N \in \mathbb{N}_0$ the completion of $C_c^\infty(\overset{\circ}{X})$ w.r.t. the norm:

$$\|v\|_{H_b^N(X)} = \left(\int_X \sum_{|I| \leq N} |\tilde{Z}^I v|^2 \frac{dvol_{m_0}}{\bar{\rho}} \right)^{\frac{1}{2}}.$$

Define the conormal symbol space by

$$\mathcal{A}^0(X) = \{v \in C^{-\infty}(X) : \mathcal{V}_b(X)^l v \in L^\infty(X), \forall l \in \mathbb{N}_0\}.$$

The relationship between the b-Sobolev spaces and the conormal symbol spaces are stated in the following lemma.

Lemma 2.2. If $c_0, c_1, c_2 > 0$, then

$$\rho_0^{c_0} \rho_1^{c_1} \rho_2^{c_2} \mathcal{A}^0(X) \subset \bigcap_{N=0}^{\infty} H_b^N(X).$$

There exists a constant C_N for any $N > \frac{1}{2}\dim(X)$ such that

$$\|v\|_{L^\infty(X)} \leq C_N \|v\|_{H_b^N(X)},$$

which implies that

$$\bigcap_{N=0}^{\infty} H_b^N(X) \subset \mathcal{A}^0(X).$$

Definition 2 and Lemma 2.2 can be generalize to any closed p -submanifolds of X with induced Riemannian metric and boundary defining functions. See [Me] for definition of p submainifold. In this paper, we mainly consider the Cauchy surface $\Sigma_0 = \{t = 0\} = \overline{\mathbb{R}^n}$ and null infinity $S_1^\pm$, which are both p -submanifolds of X . We study the map from Cauchy data to Characteristic data for Einstein vacuum equations, which lie in some weighted b-Sobolev spaces or conormal symbol space on Σ_0 and $S_1^\pm$ correspondingly.

3. INITIAL DATA

In this section, we study the spaces of small asymptotically flat initial data for Einstein vacuum equations which satisfy either the constraint equations (1.2) or the harmonic gauge conditions (1.5), and the relation between them.

Denote by $g = m + h$ a small perturbation of Minkowski metric: $g = g_{\mu\nu}dx^\mu dx^\nu$, $h = h_{\mu\nu}dx^\mu dx^\nu$. We can view g, h as functions of (t, x) valued in $(n+1) \times (n+1)$ symmetric matrices. Then (1.2) and (1.5) can be written in terms of the components of $(h^0, h^1) = (h|_{t=0}, \partial h|_{t=0})$, which turn out to be undetermined elliptic systems after suitable decomposition.

3.1. Initial Data Subject to the Constraint Equations. Let us first derive the constraint equations (1.2) in terms of components of (h^0, h^1) . Denote by

$$\Gamma_{\alpha\beta,\mu} = 2g_{\mu\nu}\Gamma_{\alpha\beta}^\nu = \partial_\alpha h_{\mu\beta} + \partial_\beta h_{\alpha\mu} - \partial_\mu h_{\alpha\beta}, \quad \Gamma_\mu = g_{\mu\nu}g^{\alpha\beta}\Gamma_{\alpha\beta}^\nu = g^{\alpha\beta}(\partial_\alpha h_{\mu\beta} - \frac{1}{2}\partial_\mu h_{\alpha\beta}).$$

Then the components of Ricci curvature of g can be expressed as

$$\begin{aligned} R_{\mu\nu} &= \partial_\lambda \Gamma_{\mu\nu}^\lambda - \partial_\mu \Gamma_{\lambda\nu}^\lambda + \Gamma_{\mu\nu}^\lambda \Gamma_{\lambda\delta}^\delta - \Gamma_{\mu\lambda}^\delta \Gamma_{\nu\delta}^\lambda \\ &= \frac{1}{2}g^{\lambda\delta}(-\partial_\lambda \partial_\delta h_{\mu\nu} + \partial_\lambda \partial_\nu h_{\mu\delta} + \partial_\lambda \partial_\mu h_{\delta\nu} - \partial_\mu \partial_\nu h_{\lambda\delta}) + E_{\mu\nu} \\ &= -\frac{1}{2}g^{\lambda\delta}\partial_\lambda \partial_\delta h_{\mu\nu} + \frac{1}{2}\partial_\mu \Gamma_\nu + \frac{1}{2}\partial_\nu \Gamma_\mu + \frac{1}{2}F_{\mu\nu} \end{aligned}$$

where

$$\begin{aligned} F_{\mu\nu} &= 2E_{\mu\nu} + D_{\mu\nu} + D_{\nu\mu}, \\ E_{\mu\nu} &= \Gamma_{\mu\nu}^\lambda \Gamma_{\lambda\delta}^\delta - \Gamma_{\mu\lambda}^\delta \Gamma_{\nu\delta}^\lambda + \frac{1}{2}(\partial_\lambda g^{\lambda\delta})\Gamma_{\mu\nu,\delta} - \frac{1}{2}(\partial_\mu g^{\lambda\delta})\Gamma_{\lambda\nu,\delta}, \\ D_{\mu\nu} &= -(\partial_\mu g^{\lambda\delta})(\partial_\lambda h_{\nu\delta} - \frac{1}{2}\partial_\nu h_{\lambda\delta}). \end{aligned}$$

Denote by

$$D'_{\mu\nu} = D_{\mu\nu}|_{t=0}, \quad E'_{\mu\nu} = E_{\mu\nu}|_{t=0}, \quad F'_{\mu\nu} = F_{\mu\nu}|_{t=0}.$$

Then $D'_{\mu\nu}, E'_{\mu\nu}, F'_{\mu\nu}$ are quadratic forms of $(\partial_i h^0, h^1)$ with coefficients analytically depending on h^0 .

The Einstein vacuum equations (1.1) imply that

$$g^{00}R_{00} - g^{ij}R_{ij} = 0, \quad g^{00}R_{0k} + g^{0j}R_{kj} = 0, \quad \text{for } k = 1, \dots, n,$$

which cancel the $\partial_0^2 h$ terms and impose $(n+1)$ constraint equations on (h^0, h^1) at $t = 0$:

$$(3.1) \quad \begin{cases} g^{pl}g^{ij}\partial_i(\partial_j h_{pl}^0 - \partial_p h_{jl}^0) + g^{pl}g^{0i}(\partial_i h_{pl}^1 - \partial_p h_{il}^1 - \partial_i \partial_p h_{0l}^0 + \partial_p \partial_l h_{i0}^0) + g^{00}E'_{00} - g^{pl}E'_{pl} = 0, \\ g^{00}g^{ij}(\partial_i h_{kj}^1 - \partial_k h_{ij}^1 - \partial_i \partial_j h_{k0}^0 + \partial_k \partial_i h_{0j}^0) + g^{0l}g^{0i}(\partial_k h_{il}^1 - \partial_i h_{kl}^1 + \partial_i \partial_l h_{k0}^0 - \partial_k \partial_l h_{0i}^0) \\ \quad + g^{0l}g^{ij}(-\partial_i \partial_j h_{kl}^0 + \partial_i \partial_k h_{jl}^0 + \partial_i \partial_l h_{kj}^0 - \partial_k \partial_l h_{ij}^0) + 2(g^{00}E'_{0k} + g^{0l}E'_{kl}) = 0. \end{cases}$$

for $k = 1, \dots, n$. Here (3.1) is the system of constraint equations (1.2) in fixed coordinates. We are interested in small solutions to (3.1) in the following space:

$$(3.2) \quad \rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \mathbb{R}^{\frac{(n+1)(n+2)}{2}}) \times \rho_0^{\lambda+1} H_b^N(\overline{\mathbb{R}^n} : \mathbb{R}^{\frac{(n+1)(n+2)}{2}}).$$

Denote by $\tilde{\mathcal{U}}_\epsilon^{N,\lambda}$ the set of solutions to (3.1) in above space with norm less than ϵ for some $\epsilon > 0$.

Proposition 3.1. *For $n \geq 3$, $\lambda \in (0, n-2)$ and $\epsilon > 0$ small enough, $\tilde{\mathcal{U}}_\epsilon^{N,\lambda}$ is a Fréchet manifold.*

Proof. We prove the statement by studying the linearization of (3.1) at $(0, 0)$:

$$(3.3) \quad \begin{cases} -\Delta \sum_i h_{ii}^0 - \sum_{ij} \partial_i \partial_j h_{ij}^0 = 0, \\ -\Delta h_{0k}^0 - \partial_k \sum_i \partial_i h_{0i}^0 + \partial_k \sum_i h_{ii}^1 - \sum_i \partial_i h_{ki}^1 = 0, \quad \text{for } k = 1, \dots, n \end{cases}$$

where $\Delta = -\sum_{j=1}^n \partial_j^2$, and applying implicit function theorem. Notice that by [Me2], for $n \geq 3$ and $\lambda \in (0, n-2)$,

$$\Delta : \rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n}) \longrightarrow \rho_0^{\lambda+2} H_b^{N-1}(\overline{\mathbb{R}^n})$$

is an isomorphism and hence

$$\Delta^{\frac{1}{2}} : \rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n}) \longrightarrow \rho_0^{\lambda+1} H_b^N(\overline{\mathbb{R}^n}) \quad \text{and} \quad \Delta^{\frac{1}{2}} : \rho_0^{\lambda+1} H_b^{N+1}(\overline{\mathbb{R}^n}) \longrightarrow \rho_0^{\lambda+2} H_b^N(\overline{\mathbb{R}^n}),$$

are both isomorphism. See Proposition 8.1 in Appendix. Assume (h^0, h^1) is in space (3.2) and take a linear transformation of as follows: for $1 \leq k \leq n$, $2 \leq l \leq n$, $1 \leq i \neq j \leq n$,

$$\begin{aligned} A_1 &= \frac{n-1}{n} \sum_i h_{ii}^0, & A_l &= \frac{1}{n} \sum_i h_{ii}^0 - h_{ll}^0, & A_{ij} &= h_{ij}^0, \\ C_1 &= \frac{n-1}{n} \sum_i h_{ii}^1, & C_l &= \frac{1}{n} \sum_i h_{ii}^1 - h_{ll}^1, & C_{ij} &= h_{ij}^1. \end{aligned}$$

Then the first equation in (3.3) becomes

$$(3.4) \quad \Delta A_1 = -\partial_1^2 \sum_{j=2}^n A_j + \sum_{j=2}^n \partial_j^2 A_j - \sum_{i \neq j} \partial_i \partial_j A_{ij}.$$

Denote by $B = h_{0k}^0 dx^k$ and $D = \Delta^{-1}(\partial_k \sum_i h_{ii}^1 - \sum_i \partial_i h_{ki}^1) dx^k$ 1-forms on the Euclidean space, i.e.

$$B, D \in \rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \Lambda^1(\mathbb{R}^n)).$$

Then the second equation in (3.3) can be expressed as

$$(3.5) \quad -\Delta B + d\delta B + \Delta D = 0,$$

Since Δ has trivial kernel on $\rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \Lambda^p(\mathbb{R}^n))$ for all $0 \leq p \leq n$, the Hodge decomposition theorem implies

$$\rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \Lambda^1(\mathbb{R}^n)) = \mathcal{E}_c^{N+1, \lambda} \oplus \mathcal{E}_{cc}^{N+1, \lambda}$$

with $\mathcal{E}_c^{N+1, \lambda}$ and $\mathcal{E}_{cc}^{N+1, \lambda}$ the subspace of closed and coclosed forms respectively. More explicitly, let

$$\begin{aligned} \Psi : \rho^{\lambda+1} H_b^N(\overline{\mathbb{R}^n}) \ni f &\longrightarrow \Delta^{-1} df \in \rho^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \Lambda^1(\mathbb{R}^n)), \\ \Phi : \rho^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \Lambda^1(\mathbb{R}^n)) \ni u &\longrightarrow \delta u \in \rho^{\lambda+1} H_b^N(\overline{\mathbb{R}^n}). \end{aligned}$$

Then $\Phi\Psi = Id$ and

$$\Psi\Phi : \rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \Lambda^1(\mathbb{R}^n)) \longrightarrow \mathcal{E}_c^{N+1, \lambda}$$

is a projection. Writing $B = B' + B''$ and $D = D' + D''$ with B', D' closed and B'', D'' coclosed, i.e.

$$B' = \Psi\Phi(B), \quad B'' = (Id - \Psi\Phi)(B), \quad D' = \Psi\Phi(D), \quad D'' = (Id - \Psi\Phi)(D),$$

then (3.5) split into

$$(3.6) \quad \Delta D' = 0, \quad -\Delta B'' + \Delta D'' = 0.$$

For the first equation in (3.6), notice that we have the following commutative diagram:

$$\begin{array}{ccc} \rho^{\lambda+1} H_b^N(\overline{\mathbb{R}^n} : \Lambda(\mathbb{R}^n)) & \xrightarrow{d, \delta} & \rho^{\lambda+2} H_b^{N-1}(\overline{\mathbb{R}^n} : \Lambda(\mathbb{R}^n)) \\ \Delta^{-\frac{1}{2}} \downarrow & & \downarrow \Delta^{-\frac{1}{2}} \\ \rho^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \Lambda(\mathbb{R}^n)) & \xrightarrow{d, \delta} & \rho^{\lambda+1} H_b^N(\overline{\mathbb{R}^n} : \Lambda(\mathbb{R}^n)) \end{array}$$

where $\Lambda(\mathbb{R}^n) = \sum_{p=0}^n \Lambda^p(\mathbb{R}^n)$. Since $h^1 \in \rho^{\lambda+1} H_b^N(\overline{\mathbb{R}^n} : \Lambda^1(\mathbb{R}^n))$,

$$\begin{aligned} \Delta D' &= \Delta \Psi\Phi(D) = d\delta \Delta^{-1}((\partial_k \sum_i h_{ii}^1 - \sum_i \partial_i h_{ki}^1) dx^k) \\ &= d\Delta^{-\frac{1}{2}} \delta((\partial_k \sum_i \Delta^{-\frac{1}{2}} h_{ii}^1 - \sum_i \partial_i \Delta^{-\frac{1}{2}} h_{ki}^1) dx^k) \\ &= d\Delta^{-\frac{1}{2}} (\Delta \sum_i \Delta^{-\frac{1}{2}} h_{ii}^1 + \sum_{ik} \partial_k \partial_i \Delta^{-\frac{1}{2}} h_{ki}^1) \end{aligned}$$

Hence $\Delta D' = 0$ is equivalent to

$$\Delta \sum_i \Delta^{-\frac{1}{2}} h_{ii}^1 + \sum_{ik} \partial_k \partial_i \Delta^{-\frac{1}{2}} h_{ki}^1 = 0.$$

Substitute h_{ij}^1 by C , (3.3) is equivalent to the elliptic system:

$$(3.7) \quad \begin{cases} \Delta A_1 = -\partial_1^2 \sum_{j=2}^n A_j + \sum_{j=2}^n \partial_j^2 A_j - \sum_{i \neq j} \partial_i \partial_j A_{ij}, \\ \Delta^{\frac{1}{2}} C_1 = -\partial_1^2 \sum_{j=2}^n \Delta^{-\frac{1}{2}} C_j + \sum_{j=2}^n \partial_j^2 \Delta^{-\frac{1}{2}} C_j - \sum_{i \neq j} \partial_i \partial_j \Delta^{-\frac{1}{2}} C_{ij}, \\ \Delta B'' = \partial_1 (C_1 - \sum_{l=2}^n C_l) dx^1 + \sum_{k=2}^n \partial_k (C_1 + C_l) dx^k + \sum_{i \neq k} \partial_i C_{ki} dx^k. \end{cases}$$

In (3.3), the components $\{h_{00}^0, h_{00}^1, h_{0k}^1 : 1 \leq k \leq n\}$ are free. To solve $\{h_{0k}^0, h_{ij}^0, h_{ij}^1 : 1 \leq i, j, k \leq n\}$ from (3.3), it is equivalent to solve $\{B_k, A_k, A_{ij}, C_i, C_{ij} : 1 \leq k, i \neq j \leq n\}$ from (3.7). Given arbitray $h_{00}^0, A_k, A_{ij} \in \rho_0^\lambda H_b^{N+1}(\mathbb{R}^n)$ and $h_{0\mu}^1, C_k, C_{ij} \in \rho_0^{\lambda+1} H_b^N(\mathbb{R}^n)$ for $0 \leq \mu \leq n, 2 \leq k \leq n, 1 \leq i \neq j \leq n$ and $B' \in \mathcal{E}_c^{N+1, \lambda}$, we can solve (3.7) uniquely and have

$$A_1 \in \rho_0^\lambda H_b^{N+1}(\mathbb{R}^n), \quad C_1 \in \rho_0^{\lambda+1} H_b^N(\mathbb{R}^n), \quad B'' \in \mathcal{E}_{cc}^{N+1, \lambda}.$$

And hence the solution space of (3.3), denoted by $\mathcal{U}^{N, \lambda}$, is isomorphism to

$$\rho_0^\lambda H_b^{N+1}(\mathbb{R}^n : \mathbb{R}^{\frac{n(n+1)}{2}}) \times \mathcal{E}_c^{N+1, \lambda} \times \rho_0^{\lambda+1} H_b^N(\mathbb{R}^n : \mathbb{R}^{\frac{(n+1)(n+2)}{2}-1})$$

Due to the ellipticity of (3.7), when $\epsilon > 0$ is small enough, $\tilde{\mathcal{U}}_\epsilon^{N, \lambda}$ is a submanifold of space (3.2), isomorphic to a small neighborhood of $\mathcal{U}^{N, \lambda}$ around $(0, 0)$ and with tangent space $\mathcal{U}^{N, \lambda}$ at $(0, 0)$. $\square$

3.2. Initial Data Subject to Harmonic Gauge Conditions. We first deriving the equations (1.5) in terms of components of (h^0, h^1) . First $\Gamma_\mu|_{t=0} = 0$ is equivalent to

$$(3.8) \quad \begin{cases} \frac{1}{2} g^{00} h_{00}^1 - \frac{1}{2} g^{ij} h_{ij}^1 + g^{i\beta} \partial_i h_{0\beta}^0 = 0, \\ g^{0\beta} h_{k\beta}^1 + g^{i\beta} \partial_i h_{k\beta}^0 - \frac{1}{2} g^{\alpha\beta} \partial_k h_{\alpha\beta}^0 = 0, \end{cases}$$

for $k = 1, \dots, n$. Secondly $\partial_t \Gamma_\mu|_{t=0}$ is equivalent to

$$(3.9) \quad \begin{cases} (\frac{1}{2} g^{00} \partial_t^2 h_{00} - \frac{1}{2} g^{ij} \partial_t^2 h_{ij})|_{t=0} + g^{i\beta} \partial_i h_{0\beta}^1 - D'_{00} = 0, \\ (g^{0\beta} \partial_t^2 h_{k\beta})|_{t=0} + g^{i\beta} \partial_i h_{k\beta}^1 - \frac{1}{2} g^{\alpha\beta} \partial_k h_{\alpha\beta}^1 - D'_{0k} = 0, \end{cases}$$

for $k = 1, \dots, n$. When assuming $\Gamma_\mu = 0$, the Reduced Einstein vacuum equations (1.3) is

$$(3.10) \quad g^{00} \partial_t^2 h_{\mu\nu}|_{t=0} = -2g^{0i} \partial_i h_{\mu\nu}^1 - g^{ij} \partial_i \partial_j h_{\mu\nu}^0 + F'_{\mu\nu}.$$

Multiplying (3.9) by g^{00} and substituting $(g^{00} \partial_t^2 h_{\mu\nu})|_{t=0}$ terms by the right hand side of (3.10), we have

$$(3.11) \quad \begin{cases} g^{00} g^{ij} \partial_i h_{0j}^1 + g^{pl} g^{0i} \partial_i h_{pl}^1 - \frac{1}{2} g^{00} g^{ij} \partial_i \partial_j h_{00}^0 + \frac{1}{2} g^{pl} g^{ij} \partial_i \partial_j h_{pl}^0 + g^{00} E'_{00} - \frac{1}{2} g^{pl} F'_{pl} = 0, \\ g^{0\beta} (-2g^{0i} \partial_i h_{k\beta}^1 - g^{ij} \partial_i \partial_j h_{k\beta}^0) + g^{00} (g^{i\beta} \partial_i h_{k\beta}^1 - \frac{1}{2} g^{\alpha\beta} \partial_k h_{\alpha\beta}^1) + g^{0\beta} F'_{k\beta} - g^{00} D'_{0k} = 0, \end{cases}$$

for $k = 1, \dots, n$.

Lemma 3.1. *The harmonic gauge conditions (3.8) and (3.11) are equivalent to the constraint conditions (3.1) combined with (3.8) in the fixed coordinates.*

Proof. We only need to show under the condition (3.8), (3.11) is equivalent to (3.1). First (3.8) gives

$$(3.12) \quad \begin{cases} g^{00} h_{00}^1 = g^{ij} h_{ij}^1 - 2g^{i\beta} \partial_i h_{0\beta}^0, \\ g^{00} h_{0k}^1 = -g^{0l} h_{kl}^1 - g^{i\beta} \partial_i h_{k\beta}^0 + \frac{1}{2} g^{\alpha\beta} \partial_k h_{\alpha\beta}^0, \end{cases}$$

for $k = 1, \dots, n$. Plugging them in the first equation of (3.11), we have

$$\begin{aligned} 0 &= g^{ij} \partial_i (-g^{0l} h_{jl}^1 - g^{l\beta} \partial_l h_{j\beta}^0 + \frac{1}{2} g^{\alpha\beta} \partial_j h_{\alpha\beta}^0) - g^{ij} \partial_i g^{00} h_{0j}^1 + g^{pl} g^{0i} \partial_i h_{pl}^1 - \frac{1}{2} g^{00} g^{ij} \partial_i \partial_j h_{00}^0 \\ &\quad + \frac{1}{2} g^{pl} g^{ij} \partial_i \partial_j h_{pl}^0 + g^{00} E'_{00} - \frac{1}{2} g^{pl} F'_{pl} \\ &= g^{0i} g^{pl} (\partial_i h_{pl}^1 - \partial_p h_{il}^1) + g^{ij} (-g^{l\beta} \partial_i \partial_l h_{j\beta}^0 + g^{pl} \partial_i \partial_j h_{pl}^0 + g^{0l} \partial_i \partial_j h_{0l}^0) + Q_0 \end{aligned}$$

where Q_0 is a quadratic form of $(\partial_i h^0, h^1)$,

$$\begin{aligned} Q_0 &= g^{ij}(-\partial_i g^{0l} h_{jl}^1 - \partial_i g^{l\beta} \partial_l h_{j\beta}^0 + \frac{1}{2} \partial_i g^{\alpha\beta} \partial_j h_{\alpha\beta}^0) - g^{ij} \partial_i g^{00} h_{0j}^1 + g^{00} E'_{00} - \frac{1}{2} g^{pl} F'_{pl} \\ &= g^{ij}(-\partial_i g^{0l} h_{jl}^1 - \partial_i g^{l\beta} \partial_l h_{j\beta}^0 + \frac{1}{2} \partial_i g^{\alpha\beta} \partial_j h_{\alpha\beta}^0 - \partial_i g^{00} h_{0j}^1 - D'_{ij}) + g^{00} E'_{00} - g^{pl} E'_{pl} \\ &= g^{00} E'_{00} - g^{pl} E'_{pl}, \end{aligned}$$

which gives the first equation of (3.1). Rewrite the second equation of (3.11) as follows:

$$\begin{aligned} & -\frac{1}{2} g^{00} g^{00} \partial_k h_{00}^1 - g^{00} g^{0i} (\partial_i h_{k0}^1 + \partial_k h_{0i}^1) + g^{00} g^{ij} (-\partial_i \partial_j h_{k0}^0 + \partial_i h_{kj}^1 - \frac{1}{2} \partial_k h_{ij}^1) \\ & - 2g^{0i} g^{0l} \partial_i h_{kl}^1 - g^{0l} g^{ij} \partial_i \partial_j h_{kl}^0 + g^{0\beta} F'_{k\beta} - g^{00} E''_{0k} = 0. \end{aligned}$$

Plugging (3.12) in the first three terms in above equation, we get

$$\begin{aligned} 0 &= -g^{00} \partial_k (\frac{1}{2} g^{ij} h_{ij}^1 - g^{i\beta} \partial_i h_{0\beta}^0) + \frac{1}{2} g^{00} \partial_k g^{00} h_{00}^1 + g^{0i} \partial_i (g^{0l} h_{kl}^1 + g^{l\beta} \partial_l h_{k\beta}^0 - \frac{1}{2} g^{\alpha\beta} \partial_k h_{\alpha\beta}^0) \\ &+ g^{0i} \partial_i g^{00} h_{k0}^1 + g^{0i} \partial_k (g^{0l} h_{il}^1 + g^{l\beta} \partial_l h_{i\beta}^0 - \frac{1}{2} g^{\alpha\beta} \partial_i h_{\alpha\beta}^0) + g^{0i} \partial_k g^{00} h_{0i}^1 \\ &+ g^{00} g^{ij} (-\partial_i \partial_j h_{k0}^0 + \partial_i h_{kj}^1 - \frac{1}{2} \partial_k h_{ij}^1) - 2g^{0i} g^{0l} \partial_i \partial_l h_{kl}^1 - g^{0l} g^{ij} \partial_i \partial_j h_{kl}^0 + g^{0\beta} F'_{k\beta} - g^{00} E''_{0k} \\ &= g^{00} g^{ij} (\partial_i h_{kj}^1 - \partial_k h_{ij}^1 - \partial_i \partial_j h_{k0}^0 + \partial_k \partial_i h_{0j}^0) + g^{0l} g^{0i} (\partial_k h_{il}^1 - \partial_i h_{kl}^1 + \partial_i \partial_l h_{k0}^0 - \partial_k \partial_l h_{0i}^0) \\ &+ g^{0l} g^{ij} (-\partial_i \partial_j h_{kl}^0 + \partial_i \partial_k h_{jl}^0 + \partial_i \partial_l h_{kj}^0 - \partial_k \partial_l h_{ij}^0) + Q_k \end{aligned}$$

where Q_k is a quadratic form of $(\partial_i h^0, h^1)$,

$$\begin{aligned} Q_k &= g^{00} (-\frac{1}{2} \partial_k g^{ij} h_{ij}^1 + \partial_k g^{i\beta} \partial_i h_{0\beta}^0 + \frac{1}{2} \partial_k g^{00} h_{00}^1) + g^{0i} (\partial_i g^{0l} h_{kl}^1 + \partial_i g^{l\beta} \partial_l h_{k\beta}^0 - \frac{1}{2} \partial_i g^{\alpha\beta} \partial_k h_{\alpha\beta}^0 \\ &+ \partial_i g^{00} h_{k0}^1 + \partial_k g^{0l} h_{il}^1 + \partial_k g^{l\beta} \partial_l h_{i\beta}^0 - \frac{1}{2} \partial_k g^{\alpha\beta} \partial_i h_{\alpha\beta}^0 + \partial_k g^{00} h_{0i}^1) + g^{0\beta} F'_{k\beta} - g^{00} E''_{0k} \\ &= -g^{00} E''_{k0} - g^{0i} (E''_{ik} + E''_{ki}) + g^{00} F'_{k0} + g^{0i} F'_{ki} - g^{00} E''_{0k} \\ &= 2(g^{00} E'_{k0} + g^{0i} E'_{ki}) \end{aligned}$$

which gives the second equation in (3.1). Since the substituting action is reversible, we finish proving the equivalence of (3.8)+(3.11) and (3.8)+(3.1). $\square$

Notice that (3.8) is an elliptic system of equations of $h_{0\mu}^1$ for $\mu = 0, 1, \dots, n$ and which is free when we solve the linearization of (3.1). Hence by a similar decomposition as in the proof of Proposition 3.1, the linearization of (3.8) + (3.11) form an elliptic system w.r.t. $\{A_1, C_1, B'', h_{0\mu}^1 : \mu = 0, \dots, n\}$ when considering the solutions in space (3.2) for or $n \geq 3$ and $\lambda \in (0, n-2)$. Denote by $\tilde{\mathcal{V}}_\epsilon^{N,\lambda}$ the space of solutions to (3.8)+(3.11) such that

$$\|(h^0, h^1)\|_{\rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n}) \times \rho_0^{\lambda+1} H_b^N(\overline{\mathbb{R}^n})} \leq \epsilon.$$

Then by the same proof of Proposition 3.1, we have

Proposition 3.2. *For $n \geq 3$, $\lambda \in (0, n-2)$ and $\epsilon > 0$ small enough, $\tilde{\mathcal{V}}_\epsilon^{N,\lambda}$ is a Fréchet manifolds.*

Here $\tilde{\mathcal{V}}_\epsilon^{N,\lambda}$ has tangent space $\mathcal{V}^{N,\lambda}$ at $(0, 0)$ which is the space of solutions to linearization of (3.8)+(3.11) and is isomorphic to

$$\rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \mathbb{R}^{\frac{n(n+1)}{2}}) \times \mathcal{E}_c^{N+1,\lambda} \times \rho_0^{\lambda+1} H_b^N(\overline{\mathbb{R}^n} : \mathbb{R}^{\frac{(n+1)n}{2}-1}).$$

3.3. Gauge Fixing. Suppose $(\mathbb{R}_{t,\bar{x}}^{1+n}, \bar{g})$ is an Einstein vacuum solution which is close to the Minkowski space-time with Cauchy data $(\bar{h}^0, \bar{h}^1)$ satisfying the constraint equations (3.1). Here

$$\bar{h}_{\mu\nu}^0 = \bar{g}_{\mu\nu} - m_{\mu\nu}|_{\bar{t}=0}, \quad \bar{h}_{\mu\nu}^1 = \partial_{\bar{t}} \bar{g}_{\mu\nu}|_{\bar{t}=0}$$

in the sense of matrices. Let $\Phi : \mathbb{R}_{t,x}^{1+n} \rightarrow \mathbb{R}_{\bar{t},\bar{x}}^{1+n}$ be a diffeomorphism preserving the Cauchy surface:

$$\bar{x}^\mu = x^\mu + f^\mu(t, x) \quad \text{with} \quad f^0(0, x) = 0.$$

We want to find out all Ψ such that (t, x) are wave coordinates w.r.t. the pull back metric $g = \Phi^* \bar{g}$:

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{g}_{\mu\beta} \partial_\nu f^\beta + \bar{g}_{\alpha\nu} \partial_\mu f^\alpha + \bar{g}_{\alpha\beta} \partial_\mu f^\alpha \partial_\nu f^\beta,$$

Denote by (h^0, h^1) the corresponding Cauchy data in coordinates (t, x) :

$$(3.13) \quad \begin{cases} h_{\mu\nu}^0 = \bar{h}_{\mu\nu}^0 + (\bar{g}_{\mu\beta}\partial_\nu f^\beta + \bar{g}_{\alpha\nu}\partial_\mu f^\alpha + \bar{g}_{\alpha\beta}\partial_\mu f^\alpha\partial_\nu f^\beta)|_{t=0}, \\ h_{\mu\nu}^1 = \bar{h}_{\mu\nu}^1 + (\bar{h}_{\mu\beta}^1\partial_\nu f^\beta + \bar{h}_{\alpha\nu}^1\partial_\mu f^\alpha + \bar{h}_{\alpha\beta}^1\partial_\mu f^\alpha\partial_\nu f^\beta)|_{t=0} \\ \quad + [\partial_t f^\gamma(\partial_{\bar{x}^\gamma}\bar{g}_{\mu\nu} + \partial_{\bar{x}^\gamma}\bar{g}_{\mu\beta}\partial_\nu f^\beta + \partial_{\bar{x}^\gamma}\bar{g}_{\alpha\nu}\partial_\mu f^\alpha + \partial_{\bar{x}^\gamma}\bar{g}_{\alpha\beta}\partial_\mu f^\alpha\partial_\nu f^\beta)]|_{t=0} \\ \quad + [\bar{g}_{\mu\beta}\partial_\nu\partial_t f^\beta + \bar{g}_{\alpha\nu}\partial_\mu\partial_t f^\alpha + \bar{g}_{\alpha\beta}(\partial_t\partial_\mu f^\alpha\partial_\nu f^\beta + \partial_\mu f^\alpha\partial_t\partial_\nu f^\beta)]|_{t=0}. \end{cases}$$

To make g satisfy the harmonic gauge condition (1.4), it is equivalent to require f satisfying the following equations:

$$(3.14) \quad (\bar{g}_{\mu\nu} + \bar{g}_{\delta\nu}\partial_\mu f^\delta)g^{\alpha\beta}\partial_\alpha\partial_\beta f^\nu + G_\mu(\bar{g}, \partial\bar{g}; f, \partial f) = 0, \quad \text{for } \mu = 0, \dots, n,$$

where G_μ are analytic functions of $(\bar{g}, \partial\bar{g}; f, \partial f)$:

$$\begin{aligned} G_\mu(\bar{g}, \partial\bar{g}; f, \partial f) &= g^{\alpha\beta}(\partial_\alpha\bar{g}_{\mu\beta} + \partial_\alpha\bar{g}_{\mu\gamma}\partial_\beta f^\gamma + \partial_\alpha\bar{g}_{\delta\beta}\partial_\mu f^\delta + \bar{g}_{\delta\gamma}\partial_\mu f^\delta\partial_\beta f^\gamma) \\ &\quad - \frac{1}{2}g^{\alpha\beta}(\partial_\mu\bar{g}_{\alpha\beta} + \partial_\mu\bar{g}_{\alpha\gamma}\partial_\beta f^\gamma + \partial_\mu\bar{g}_{\delta\beta}\partial_\alpha f^\delta + \partial_\mu\bar{g}_{\delta\gamma}\partial_\alpha f^\delta\partial_\beta f^\gamma), \\ G_\mu(\bar{g}, \partial\bar{g}; 0, 0) &= \bar{g}^{\alpha\beta}(\bar{\partial}_\alpha\bar{g}_{\mu\beta} - \frac{1}{2}\bar{\partial}_\mu\bar{g}_{\alpha\beta}). \end{aligned}$$

Notice that $\partial_\alpha\bar{g}_{\mu\nu} = \bar{\partial}_\alpha\bar{g}_{\mu\nu} + \partial_\alpha f^\beta\bar{\partial}_\beta\bar{g}_{\mu\nu}$. Here (3.14) is a system of quasilinear wave equations of f . Given any $f_0 = f|_{t=0}$ and $f_1 = \partial_t f|_{t=0}$ with enough regularity, we can solve (3.14) in a neighborhood of Cauchy surface. Furthermore, if (f_0, f_1) are also small enough, f is determined globally. Since we assume $f_0^0 = 0$, (f_0, f_1) have $2n + 1$ freedom.

However, we want to choose the gauge before we find out the global Einstein solution $\bar{g}$ for Cauchy problem. By Theorem 1, this can be done by requiring Cauchy data (h^0, h^1) in the new coordinates (t, x) to satisfy the harmonic gauge conditions (1.5), or equivalently, (3.8) and (3.11). First (3.8) give $(n + 1)$ linear equations of $\partial_t^2 f|_{t=0}$ and allow us to write $\partial_t^2 f|_{t=0}$ in terms of (f_0, f_1) . More explicitly

$$(3.15) \quad \partial_t^2 f^\mu|_{t=0} = f_2^\mu(\bar{h}^0, \bar{h}^1; f_0, f_1)$$

where f_2^μ are analytic functions of $(\bar{h}^0, \bar{h}^1; f_0, f_1)$. Then (3.11) is satisfied automatically by Lemma 3.1 since the constraint equations (1.2), or equivalently (3.1), is invariant under diffeomorphism. This shows again that the global diffeomorphism, or equivalently the global harmonic coordinates set, is determined by (f_0, f_1) .

3.4. Group Action and Fibration. From above discussion, the set of $(f_0, f_1) = (f|_{t=0}, \partial_t f|_{t=0})$ gives a group action on the space of Cauchy data (h^0, h^1) which satisfies the harmonic gauge conditions (3.8) and (3.11) via (3.13). We discuss this group action in the following. For $\lambda > 0$, let $\mathcal{G}^{N, \lambda}$ be the nonabelian group generated by

$$\{(f_0, f_1) \in \rho_0^{\lambda-1}H_b^{N+2}(\overline{\mathbb{R}^n} : \mathbb{R}^{n+1}) \times \rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n} : \mathbb{R}^{n+1}) : f_0^0 = 0, \|(f_0, f_1)\| < \epsilon'\},$$

for some $\epsilon' > 0$ small with group operation

$$(\tilde{f}_0, \tilde{f}_1) \star (f_0, f_1) = (\tilde{f}_0 + f_0 \circ (Id + \tilde{f}_0), \tilde{f}_1 + \tilde{f}_1^0 f_1 \circ (Id + \tilde{f}_0) + \tilde{f}_1^i \partial_i f_0 \circ (Id + \tilde{f}_0)).$$

and identity $(0, 0)$. Here $\circ$ denotes the composition of functions. It is clearly that $\mathcal{G}^{N, \lambda}$ does not depend on $\epsilon' > 0$ small. Then $\mathcal{G}^{N, \lambda}$ has two subgroups generated by $(f_0, 0)$ and $(0, f_1)$ respectively, which transversely intersect at $(0, 0)$. Denote by

$$\mathcal{G}_0^{N, \lambda} = \{f_0 : (f_0, 0) \in \mathcal{G}^{N, \lambda}\}, \quad \mathcal{G}_1^{N, \lambda} = \{f_1 : (0, f_1) \in \mathcal{G}^{N, \lambda}\}.$$

Lemma 3.2. *With above notation,*

$$\mathcal{G}^{N, \lambda} = \mathcal{G}_1^{N, \lambda} \star \mathcal{G}_0^{N, \lambda} = \mathcal{G}_0^{N, \lambda} \star \mathcal{G}_1^{N, \lambda}.$$

Denote by $\tilde{\mathcal{V}}^{N, \lambda} = \mathcal{G}^{N, \lambda} \cdot \tilde{\mathcal{V}}_\epsilon^{N, \lambda} \subset \tilde{\mathcal{V}}_\infty^{N, \lambda}$ where ϵ is chosen in Proposition 3.2. By Lemma 3.2, we can study $\mathcal{G}_1^{N, \lambda}$ and $\mathcal{G}_0^{N, \lambda}$ separately. First, $\mathcal{G}_1^{N, \lambda}$ forms an nonabelian group with operation

$$(3.16) \quad (0, \tilde{f}_1) \star (0, f_1) = (0, f_1 + \tilde{f}_1 + \tilde{f}_1^0 f_1), \quad \forall f_1, \tilde{f}_1 \in \mathcal{G}_1^{N, \lambda}.$$

By (3.13), given $f_1 \in \mathcal{G}_1^{N,\lambda}$ and $(\bar{h}^0, \bar{h}^1) \in \tilde{\mathcal{V}}^{N,\lambda}$, then $(h^0, h^1) = (0, f_1) \cdot (\bar{h}^0, \bar{h}^1)$ is defined as follow:

$$(3.17) \quad \begin{cases} h_{00}^0 = \bar{h}_{00}^0 + 2\bar{g}_{0\alpha}f_1^\alpha + \bar{g}_{\alpha\beta}f_1^\alpha f_1^\beta, \\ h_{0i}^0 = \bar{h}_{0i}^0 + \bar{g}_{i\beta}f_1^\beta, \\ h_{ij}^0 = \bar{h}_{ij}^0 \\ h_{00}^1 = (\bar{h}_{00}^1 + \bar{h}_{00}^1 f_1^0 + \partial_i \bar{h}_{00}^0 f_1^i) + 2(\bar{h}_{0\beta}^1 + \bar{h}_{0\beta}^1 f_1^0 + \partial_i \bar{h}_{0\beta}^0 f_1^i) f_1^\beta \\ \quad + (\bar{h}_{\alpha\beta}^1 + \bar{h}_{\alpha\beta}^1 f_1^0 + \partial_i \bar{h}_{\alpha\beta}^0 f_1^i) f_1^\alpha f_1^\beta + 2\bar{g}_{0\beta}f_2^\beta + 2\bar{g}_{\alpha\beta}f_1^\alpha f_2^\beta, \\ h_{0i}^1 = (\bar{h}_{0i}^1 + \bar{h}_{0i}^1 f_1^0 + \partial_k \bar{h}_{0i}^0 f_1^k) + (\bar{h}_{i\beta}^1 + \bar{h}_{i\beta}^1 f_1^0 + \partial_k \bar{h}_{i\beta}^0 f_1^k) f_1^\beta \\ \quad + \bar{g}_{i\beta}f_2^\beta + \bar{g}_{0\beta}\partial_i f_1^\beta + \bar{g}_{\alpha\beta}f_1^\alpha \partial_i f_1^\beta, \\ h_{ij}^1 = \bar{h}_{ij}^1 + \bar{h}_{ij}^1 f_1^0 + \partial_k \bar{h}_{ij}^0 f_1^k + \bar{g}_{i\beta}\partial_j f_1^\beta + \bar{g}_{\alpha\beta}\partial_i f_1^\alpha. \end{cases}$$

Here $f_2 = f_2(\bar{h}^0, \bar{h}^1; 0, f_1)$ is defined in (3.15). Obviously $\mathcal{G}_1^{N,\lambda}$ preserves the coordinates on the Cauchy surface and hence preserves the restricting metric and second fundamental form on Cauchy surface. Given initial data (h_0, h_1) , let g_0 be the induced metric on Cauchy surface and k_0 the corresponding second fundamental form. This defines the map $\pi : (h^0, h^1) \rightarrow (g_0, k_0)$:

$$[g_0]_{ij} = \delta_{ij} + h_{ij}^0, \quad [k_0]_{ij} = -(1 - h_{00}^0 + g_0^{kl}h_{0l}^0 h_{0k}^0)^{\frac{1}{2}}(\partial_i h_{0j}^0 + \partial_j h_{0i}^0 - h_{ij}^1).$$

In our setting,

$$[g_0]_{ij} - \delta_{ij} \in \rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n}), \quad [k_0]_{ij} \in \rho_0^{\lambda+1} H_b^N(\overline{\mathbb{R}^n}).$$

Let $\tilde{\mathcal{C}}_\epsilon^{N,\lambda} = \pi(\mathcal{G}_1^{N,\lambda} \cdot \tilde{\mathcal{V}}_\epsilon^{N,\lambda})$. Obviously,

$$\begin{array}{ccc} \mathcal{G}_1^{N,\lambda} & \xrightarrow{\quad} & \mathcal{G}_1^{N,\lambda} \cdot \tilde{\mathcal{V}}_\epsilon^{N,\lambda} \\ & & \downarrow \pi \\ & & \tilde{\mathcal{C}}_\epsilon^{N,\lambda} \end{array}$$

provides a fibration structure.

Secondly, $\mathcal{G}_0^{N,\lambda}$ is obvious equivalent to the diffeomorphism group on Cauchy surface with the corresponding regularity and decay at infinity. The group operation is as follows:

$$\tilde{f}_0 \star f_0 = \tilde{f}_0 + f_0 \circ (Id + \tilde{f}_0), \quad \forall \tilde{f}_0, f_0 \in \mathcal{G}_0^{N,\lambda}.$$

The action of $\mathcal{G}_0^{N,\lambda}$ on $\tilde{\mathcal{C}}_\epsilon^{N,\lambda}$ is just a pull-back. Denote by $\tilde{\mathcal{C}}^{N,\lambda} = \mathcal{G}_0^{N,\lambda} \cdot \tilde{\mathcal{C}}_\epsilon^{N,\lambda}$. Then for any $f_0 \in \mathcal{G}_0^{N,\lambda}$, we have the commutativity

$$\pi \circ f_0 = f_0 \circ \pi$$

which implies that

$$\begin{array}{ccc} \mathcal{G}_1^{N,\lambda} & \xrightarrow{\quad} & \tilde{\mathcal{V}}^{N,\lambda} \\ & & \downarrow \pi \\ & & \tilde{\mathcal{C}}^{N,\lambda} \end{array}$$

is also a fibration. Let $\tilde{\mathcal{D}}^{N,\lambda} = \tilde{\mathcal{C}}^{N,\lambda} / \mathcal{G}_0^{N,\lambda}$ and π' the quotient map. Denote by m_e be the Euclidean metric and $S^2(T^*\mathbb{R}^n)$ be the symmetric 2-tensor bundle on $\mathbb{R}^n$. Summarizing above discussion, we have

Proposition 3.3. *For $n \geq 3$ and $\lambda \in (0, n-2)$, the quotient space $\tilde{\mathcal{D}}^{N,\lambda}$ is an open neighborhood of $(m_e, 0)$ in the solution space of constraint equation (1.2) in*

$$(m_e + \rho_0^\lambda H_b^{N+1}(\overline{\mathbb{R}^n}; S^2(T^*\mathbb{R}^n))) \times \rho_0^{\lambda+1} H_b^N(\overline{\mathbb{R}^n}; S^2(T^*\mathbb{R}^n)).$$

with the fibration structure:

$$\begin{array}{ccc} \mathcal{G}_1^{N,\lambda} & \xrightarrow{\quad} & \tilde{\mathcal{V}}^{N,\lambda} \\ & & \downarrow \pi \\ \mathcal{G}_0^{N,\lambda} & \xrightarrow{\quad} & \tilde{\mathcal{C}}^{N,\lambda} \\ & & \downarrow \pi' \\ & & \tilde{D}^{N,\lambda}. \end{array}$$

4. CONFORMAL TRANSFORMATION

To study the asymptotic behavior of solutions which are close to Minkowski space-time, it is equivalent to study the asymptotic behavior of their conformal transformation on X , the compactification of $\mathbb{R}_{t,x}^{1+n}$ defined in Section 2, by performing energy estimates for a conformal transformation of the reduced Einstein vacuum equation (1.3) with Cauchy data satisfying the harmonic gauge condition (3.8) and (3.11). We clarify the conformal transformation of metrics and equations in this section by describing them in the local coordinates on X specified in Section 2.

4.1. Perturbed Lorentzian Metric. Let $g = m + h$ be a Lorentzian metric on $\mathbb{R}_{t,x}^{1+n}$ with $m = -dt^2 + \sum_{x=1}^n d^2x^i$ the Minkowski metric and h a small symmetric $(0,2)$ -tensor: $h = h_{00}d^2t + 2h_{0i}dtdx^i + h_{ij}dx^i dx^j$. In the fixed coordinates $(t, x) = (x^0, x^1, \dots, x^n)$, it is equivalent to say that h is a function on $\mathbb{R}_{t,x}^{1+n}$ or the interior of X valued in $(n+1) \times (n+1)$ symmetric matrix space $SM(n+1, \mathbb{R}) \simeq \mathbb{R}^{\frac{(n+2)(n+1)}{2}}$. In this sense, denote by

$$g^{\mu\nu} = m^{\mu\nu} + H^{\mu\nu}, \quad \mu, \nu = 0, \dots, n.$$

Definition 3. For any $k \in \mathbb{N}_0$, denote by $\Theta_k(h)$ a real analytic function of h with coefficients in $\mathcal{A}^{0,0,0}(X)$ such that

$$\Theta_k(h) = O(|h|^k) \quad \text{as } |h| \rightarrow 0.$$

For any $k \in \mathbb{N}_0, j, l \in \mathbb{N}$, denote by $\Theta_k(h)(v^1, \dots, v^l)$ an l -form in $(v^1, \dots, v^l)$ with coefficients $\Theta_k(h)$, and $\Theta_k(h)(v^1, \dots, v^l)(u^1, \dots, u^j)$ an j -form in $(u^1, \dots, u^j)$ with coefficients $\Theta_k(h)(v^1, \dots, v^l)$.

Lemma 4.1. For any $k, l \in \mathbb{N}_0$

$$\Theta_k(h) + \Theta_l(h) = \Theta_{\min\{k,l\}}(h), \quad \Theta_k(h)\Theta_l(h) = \Theta_{k+l}(h).$$

Lemma 4.2. With above notation

$$H^{00} = -h_{00} + \Theta_2(h), \quad H^{0i} = h_{0i} + \Theta_2(h), \quad H^{ij} = -h_{ij} + \Theta_2(h).$$

For the purpose of studying the geometry of g near null infinity of $(\mathbb{R}_{t,x}^{1+n}, g)$, we also denote by

$$h_{0\rho} = h_{\rho 0} = h_{0\mu}\theta^\mu, \quad h_{\rho\rho} = h_{\mu\nu}\theta^\mu\theta^\nu, \quad h_{i\rho} = h_{\rho i} = h_{i\mu}\theta^\mu.$$

With this notations, the harmonic gauge condition says that some components decay faster than the others near null infinity.

Lemma 4.3. Suppose (t, x) are harmonic coordinates w.r.t. g . Then in each domain Ω_i for $1 \leq i \leq 5$ with the choice of $\tilde{\rho}$ and ρ_1 specified in Section 2, there exists $D \in \mathcal{V}_b(\Omega_i)$ such that

$$[D, \tilde{\rho}] = [D, \theta] = 0 \quad \text{and} \quad D(2h_{\mu\rho} - \theta_\mu \text{tr}_m h) = \rho_1 \tilde{\partial} h + \Theta_1(h)(\tilde{\partial} h), \quad \text{for } \mu = 0, 1, \dots, n.$$

Here $\tilde{\partial} \in \mathcal{V}_b(X)$.

Proof. We prove the statement by writing out a particular choice of D in local coordinates. First in Ω_1 and Ω_5 , it is obviously true since $\rho_1 = 1$ and we can choose $D = 0$. For the other domains, D can be chosen as follows:

- In Ω_2 , $a = 1 - \frac{t}{r}$, $b = \frac{1}{r-t}$, $\tilde{\rho} = ab$, $\rho_1 = a$,

$$\begin{aligned} (b\partial_b - a\partial_a)(2h_{0\rho} + \text{tr}_m h) &= a\tilde{\partial}h + \Theta_1(h)(\tilde{\partial}h), \\ (b\partial_b - a\partial_a)(2h_{i\rho} - \theta_i \text{tr}_m h) &= a\tilde{\partial}h + \Theta_1(h)(\tilde{\partial}h), \\ \implies Dh_{\rho\rho} &= a\tilde{\partial}h + \Theta_1(h)(\tilde{\partial}h), \quad D = (b\partial_b - a\partial_a). \end{aligned}$$

- In Ω_3 , $\rho = \frac{1}{r}$, $\tau = t - r$, $\tilde{\rho} = \rho_1 = \rho$,

$$\begin{aligned} \partial_\tau(2h_{0\rho} + \text{tr}_m h) &= \rho\tilde{\partial}h + \Theta_1(h)(\tilde{\partial}h), \\ \partial_\tau(2h_{i\rho} - \theta_i \text{tr}_m h) &= \rho\tilde{\partial}h + \Theta_1(h)(\tilde{\partial}h), \\ \implies Dh_{\rho\rho} &= \rho\tilde{\partial}h + \Theta_1(h)(\tilde{\partial}h), \quad D = \partial_\tau. \end{aligned}$$

- In Ω_4 , $\bar{a} = 1 - \frac{r}{t}$, $\bar{b} = \frac{1}{t-r}$, $\tilde{\rho} = \bar{a}\bar{b}$, $\rho_1 = \bar{a}$,

$$\begin{aligned} (\bar{a}\partial_{\bar{a}} - \bar{b}\partial_{\bar{b}})(2h_{0\rho} + \text{tr}_m h) &= \bar{a}\tilde{\partial}h + \Theta_1(h)(\tilde{\partial}h), \\ (\bar{a}\partial_{\bar{a}} - \bar{b}\partial_{\bar{b}})(2h_{i\rho} - \theta_i \text{tr}_m h) &= \bar{a}\tilde{\partial}h + \Theta_1(h)(\tilde{\partial}h), \\ \implies Dh_{\rho\rho} &= \bar{a}\tilde{\partial}h + \Theta_1(h)(\tilde{\partial}h), \quad D = (\bar{a}\partial_{\bar{a}} - \bar{b}\partial_{\bar{b}}). \end{aligned}$$

Here $\text{tr}_m h = m^{\alpha\beta} h_{\alpha\beta} = -h_{00} + \sum_{i=1}^n h_{ii}$. We finish the proof. $\square$

4.2. Conformal Transformation of the Lorentzian Metric. Recall that $\tilde{\rho}$ denotes a total boundary defining function of X and ρ_i denotes the defining function for boundary hypersurfaces S_i for $i = 0, 1, 2$. Set

$$(4.1) \quad \tilde{m} = \tilde{\rho}^2 m, \quad \tilde{h} = \tilde{\rho}^{\frac{1-n}{2}} h, \quad \tilde{g} = \tilde{\rho}^2 g = \tilde{m} + \tilde{\rho}^2 h = \tilde{m} + \tilde{\rho}^{\frac{n+3}{2}} \tilde{h}.$$

Similarly, we denote by

$$\tilde{h}_{0\rho} = \tilde{h}_{\tau 0} = \tilde{h}_{0\mu} \theta^\mu, \quad \tilde{h}_{\rho\rho} = \tilde{h}_{\mu\nu} \theta^\mu \theta^\nu, \quad \tilde{h}_{i\rho} = \tilde{h}_{\rho i} = \tilde{h}_{\mu i} \theta^\mu.$$

Lemma 4.4. *Under the assumption and notation in Lemma 4.3,*

$$D(2\tilde{h}_{\mu\rho} - \theta_\mu \text{tr}_m \tilde{h}) = \rho_1 \tilde{\partial} \tilde{h} + \rho_1 \Theta_1(\tilde{h}) + \tilde{\rho}^{\frac{n-1}{2}} (\Theta_1(\tilde{h})(\tilde{\partial} \tilde{h}) + \Theta_2(\tilde{h})), \quad \text{for } \mu = 0, 1, \dots, n.$$

Proposition 4.1. *If $h \in C^0(X : SM(n+1, \mathbb{R}))$ such that for some $\delta > 0$ and all $1 \leq i \leq n$*

$$\begin{aligned} \rho_1^{-1} h_{\rho i} &\in C^0(X : SM(n+1, \mathbb{R})), \quad \rho_1^{-1-\delta} h_{\rho\rho} \in C^0(X : SM(n+1, \mathbb{R})), \\ \sum_{\mu, \nu} \|h_{\mu\nu}\|_\infty + \sum_{1 \leq i \leq n} \|\rho_1^{-1} h_{\rho i}\|_\infty + \|\rho_1^{-1-\delta} h_{\rho\rho}\|_\infty &< \epsilon, \end{aligned}$$

for some $\epsilon > 0$ small, then $\tilde{g}$ extends to a $C^0(X^2)$ Lorentzian metric of signature $(n, 1)$ on X^2 with $S^\pm$ being its characteristic surfaces.

Proof. The statement is independent of choice of boundary defining functions. Hence we only need to write the metric $\tilde{g} = \tilde{m} + \tilde{\rho}^2 h$ in local coordinates near ∂X with particular choice of $\tilde{\rho}$ and ρ_i as stated in Section 2.

- In Ω_1 , $s = \frac{t}{r}$, $\rho = \frac{1}{r}$, $\tilde{\rho} = \rho$,

$$\begin{aligned} \tilde{m} &= -d^2 s + 2s ds \frac{d\rho}{\rho} + (1-s^2) \left(\frac{d\rho}{\rho}\right)^2 + d^2 \theta, \\ \tilde{\rho}^2 h &= h_{00} d^2 s + (h_{00} s^2 + 2h_{0i} \theta^i s + h_{ij} \theta^i \theta^j) \left(\frac{d\rho}{\rho}\right)^2 - 2(h_{00} s + h_{0i} \theta^i) ds \frac{d\rho}{\rho} \\ &\quad + h_{ij} d\theta^i d\theta^j + 2h_{0i} ds d\theta^i - 2(h_{0i} s + h_{ij} \theta^j) d\theta^j \frac{d\rho}{\rho}. \end{aligned}$$

- In Ω_2 , $a = 1 - \frac{t}{r}$, $b = \frac{1}{r-t}$, $\tilde{\rho} = ab$, $\rho_1 = a$,

$$\begin{aligned}\tilde{m} &= 2da\frac{db}{b} + a(2-a)\left(\frac{db}{b}\right)^2 + d^2\theta, \quad \tilde{m}|_{\rho_1=0} = d^2\theta, \\ \tilde{\rho}^2 h &= h_{\rho\rho}\left(\frac{da}{a}\right)^2 + (h_{\rho\rho} - 2ah_{0\rho} + a^2h_{00})\left(\frac{db}{b}\right)^2 + 2(h_{\rho\rho} - ah_{0\rho})\frac{da}{a}\frac{db}{b} \\ &\quad + h_{ij}d\theta^i d\theta^j + 2h_{\rho i}\frac{da}{a}d\theta^i + 2(-h_{\rho i} + ah_{0i})\frac{db}{b}d\theta^i.\end{aligned}$$

- In Ω_3 , $\rho = \frac{1}{r}$, $\tau = t - r$, $\tilde{\rho} = \rho_1 = \rho$,

$$\begin{aligned}\tilde{m} &= 2d\tau d\rho - \rho^2 d^2\tau + d^2\theta, \quad \tilde{m}|_{\rho_1=0} = d^2\theta, \\ \tilde{\rho}^2 h &= h_{\rho\rho}\left(\frac{d\rho}{\rho}\right)^2 + \rho^2 h_{00}d^2\tau - 2h_{0\rho}d\tau d\rho + h_{ij}d\theta^i d\theta^j + 2\rho h_{0i}d\theta^i d\tau - 2h_{\rho i}d\theta^i \frac{d\rho}{\rho}.\end{aligned}$$

- In Ω_4 , $\bar{a} = 1 - \frac{r}{t}$, $\bar{b} = \frac{1}{t-r}$, $\tilde{\rho} = \bar{a}\bar{b}$, $\rho_1 = \bar{a}$,

$$\begin{aligned}\tilde{m} &= -2d\bar{a}\frac{d\bar{b}}{\bar{b}} - \bar{a}(2-\bar{a})\left(\frac{d\bar{b}}{\bar{b}}\right)^2 + (1-\bar{a})^2 d^2\theta, \quad \tilde{m}|_{\rho_1=0} = d^2\theta, \\ \tilde{\rho}^2 h &= h_{\rho\rho}\left(\frac{d\bar{a}}{\bar{a}}\right)^2 + (h_{\rho\rho} - 2\bar{a}h_{i\rho}\theta^i + \bar{a}^2 h_{ij}\theta^i\theta^j)\left(\frac{d\bar{b}}{\bar{b}}\right)^2 + 2(h_{\rho\rho} - \bar{a}h_{\rho i}\theta^i)\frac{d\bar{a}}{\bar{a}}\frac{d\bar{b}}{\bar{b}} \\ &\quad + (1-\bar{a})^2 h_{ij}d\theta^i d\theta^j - 2(1-\bar{a})h_{\rho i}d\theta^i \frac{d\bar{a}}{\bar{a}} + 2(1-\bar{a})(-h_{\rho i} + \bar{a}h_{ij}\theta^j)d\theta^i \frac{d\bar{b}}{\bar{b}}.\end{aligned}$$

- In Ω_5 , $\phi = \frac{1}{t}$, $y = \frac{x}{t}$, $\tilde{\rho} = \phi$,

$$\begin{aligned}\tilde{m} &= -(1-|y|^2)\left(\frac{d\phi}{\phi}\right)^2 + \sum_{i=1}^n d^2 y^i - 2y^i dy^i \left(\frac{d\phi}{\phi}\right), \\ \tilde{\rho}^2 h &= (h_{00} + 2h_{0i}y^i + h_{ij}y^i y^j)\left(\frac{d\phi}{\phi}\right)^2 + h_{ij}dy^i dy^j - 2(h_{0i} + h_{ij}y^j)dy^i \frac{d\phi}{\phi}.\end{aligned}$$

Here in Ω_i for $1 \leq i \leq 4$ we use the polar coordinates. Notice here θ is restricted on $\mathbb{S}^{n-1}$, i.e. $|\theta| = 1$ and $\theta_i d\theta^i = 0$. Then the statement is obviously true if $\delta \geq 1$. For $\delta \in (0, 1)$, let

$$f(\rho_1) = \frac{1}{2} \int_0^{\rho_1} \rho_1'^{-2} h_{\rho\rho} d\rho_1' \in \rho_1^\delta C^0(X).$$

and change the coordinates near S_1 as follows:

- In Ω_2 , set $b = b'e^{-f(a)}$;
- In Ω_3 , set $\tau = \tau' - f(\rho)$;
- In Ω_4 , set $\bar{b} = \log \bar{b}' e^{f(\bar{a})}$.

In the new coordinates the metric components is uniformly bounded and $\{\rho_1 = 0\}$ is its characteristic surface. Since the coordinates changing preserves the boundary hypersurface $S_1^\pm$, we prove the statement. Notice that the coordinates change extends to a $C^{0,\delta}$ homeomorphism of $X \rightarrow X$. $\square$

4.3. Wave Operator and Commutators. Denote by $\square_g$ (resp. $\square_{\tilde{g}}$) the Laplace-Beltrami operator w.r.t. g (resp. $\tilde{g} = \tilde{\rho}^2 g$). The relation between $\square_g$ and $\square_{\tilde{g}}$ is stated as follows.

Lemma 4.5. *For any $u \in C^2(\mathbb{R}_{t,x}^{1+n})$,*

$$(4.2) \quad \square_g(\tilde{\rho}^{\frac{n-1}{2}} u) = \tilde{\rho}^{\frac{n+3}{2}} (\square_{\tilde{g}} + \gamma)u, \quad \gamma = -\tilde{\rho}^{\frac{n-1}{2}} \square_{\tilde{g}} \tilde{\rho}^{\frac{1-n}{2}}.$$

Proof. The formula can be proved by direct computation as Friedlander did in [Fr].

$$\begin{aligned}(4.3) \quad \square_g(\tilde{\rho}^{\frac{n-1}{2}} u) &= |g|^{-\frac{1}{2}} \partial_\mu (|g|^{\frac{1}{2}} g^{\mu\nu} \partial_\nu (\tilde{\rho}^{\frac{n-1}{2}} u)) \\ &= \tilde{\rho}^{n+1} |\tilde{g}|^{-\frac{1}{2}} \partial_\mu (\tilde{\rho}^{1-n} |\tilde{g}|^{\frac{1}{2}} g^{\mu\nu} \partial_\nu (\tilde{\rho}^{\frac{n-1}{2}} u)) \\ &= \tilde{\rho}^{n+1} |\tilde{g}|^{-\frac{1}{2}} \partial_\mu (|\tilde{g}|^{\frac{1}{2}} g^{\mu\nu} (\tilde{\rho}^{\frac{1-n}{2}} \partial_\nu u - u \partial_\nu \tilde{\rho}^{\frac{1-n}{2}})) \\ &= \tilde{\rho}^{\frac{n+3}{2}} (\square_{\tilde{g}} u + \gamma u)\end{aligned}$$

$\square$

Lemma 4.6. Suppose (t, x) are harmonic coordinates w.r.t. g . Then in each domain Ω_i for $1 \leq i \leq 5$, with the choice of $\tilde{\rho}, \rho_0, \rho_1, \rho_2$ specified in Section 2, we have

$$\begin{aligned}\square_{\tilde{g}} &= \square_{\tilde{m}} + (\rho_0 \rho_2)^{\frac{n-1}{2}} [-\rho_1^{\frac{n-5}{2}} \tilde{h}_{\rho\rho} (D^2 + (3-i)D) + \rho_1^{\frac{n-3}{2}} \Theta_1(\tilde{h})((\tilde{\partial}^2 + \tilde{\partial}))], \\ \gamma &= \gamma_0 + \tilde{\rho}^{\frac{n-1}{2}} \Theta_1(\tilde{h}),\end{aligned}$$

where $D \in \mathcal{V}_b(\Omega_i)$ is defined in Lemma 4.3 and $\tilde{\partial} \in \mathcal{V}_b(X)$, $\tilde{\partial}^2 \in \mathcal{V}_b^2(X)$. Here $\gamma_0 = -\tilde{\rho}^{\frac{n-1}{2}} \square_{\tilde{m}} \tilde{\rho}^{\frac{1-n}{2}}$ is a constant in each domain and in particular,

$$\square_{\tilde{m}} = 2\partial_{\rho_1} D + \tilde{\partial}^2 + \tilde{\partial} \quad \text{in } \Omega_2, \Omega_3 \text{ and } \Omega_4.$$

Proof. If (t, x) are harmonic coordinates w.r.t. g , then $\square_g = g^{\mu\nu} \partial_\mu \partial_\nu = \square_m + H^{\mu\nu} \partial_\mu \partial_\nu$, which implies that

$$\square_{\tilde{g}} + \gamma = \square_{\tilde{m}} + \gamma_0 + \tilde{\rho}^{-\frac{n+3}{2}} H^{\mu\nu} \partial_\mu \partial_\nu \tilde{\rho}^{\frac{n-1}{2}}.$$

We can write $\square_{\tilde{m}}, \tilde{\rho}^{-\frac{n+3}{2}} H^{\mu\nu} \partial_\mu \partial_\nu \tilde{\rho}^{\frac{n-1}{2}}, \gamma_0$ and γ in local coordinates in each domain Ω_i for $1 \leq i \leq 5$ with a particular choice of $\tilde{\rho}, \rho_0, \rho_1, \rho_2$ specified in Section 2.

- In Ω_1 , $s = \frac{t}{r}$, $\rho = \frac{1}{r}$, $\tilde{\rho} = \rho$,

$$\begin{aligned}\square_{\tilde{m}} &= -(1-s^2)\partial_s^2 + 2s\partial_s \rho \partial_\rho + 2s\partial_s + (\rho \partial_\rho)^2 + \rho \partial_\rho + \triangle_\theta, \\ \square_{\tilde{g}} &= \square_{\tilde{m}} + [H^{00} - 2sH^{0i}\theta_i + s^2 H^{ij}\theta_i \theta_j] \partial_s^2 + [H^{ij}\theta_i \theta_j] (\rho \partial_\rho)^2 + H^{ij} \tilde{\phi}_i \tilde{\phi}_j \\ &\quad + 2[-H^{0i}\theta_i + sH^{ij}\theta_i \theta_j] \partial_s (\rho \partial_\rho) + 2[H^{0i} - sH^{ij}\theta_j] \partial_s \tilde{\phi}_i \\ &\quad + 2[-H^{ij}\theta_j] (\rho \partial_\rho) \tilde{\phi}_i + [-(n+1)H^{0i}\theta_i + sH^{ij}(-\delta_{ij} + (n+2)\theta_i \theta_j)] \partial_s \\ &\quad + [H^{ij}(-\delta_{ij} + (n+1)\theta_i \theta_j)] \rho \partial_\rho + [-nH^{ij}\theta_j] \tilde{\phi}_i \\ &= \square_{\tilde{m}} + \rho^{\frac{n-1}{2}} \Theta_1(\tilde{h}) (\tilde{\partial}^2 + \tilde{\partial}), \\ \gamma &= \gamma_0 + [H^{ij}(-\frac{n-1}{2}\delta_{ij} + \frac{(n-1)(n+3)}{4}\theta_i \theta_j)], \quad \gamma_0 = -\frac{(n-1)(n-3)}{4}.\end{aligned}$$

- In Ω_2 , $a = 1 - \frac{t}{r}$, $b = \frac{1}{r-t}$, $\tilde{\rho} = ab$,

$$\begin{aligned}\square_{\tilde{m}} &= 2\partial_a b \partial_b - a(2-a)\partial_a^2 - 2(1-a)\partial_a + \triangle_\theta \\ &= 2\partial_a (b\partial_b - a\partial_a) + (a\partial_a)^2 + a\partial_a + \triangle_\theta \\ \square_{\tilde{g}} &= \square_{\tilde{m}} + a^{-2} [H^{00} - 2H^{i0}\theta_i + H^{ij}\theta_i \theta_j] (b\partial_b - a\partial_a)^2 + [H^{ij}\theta_i \theta_j] (a\partial_a)^2 \\ &\quad + 2a^{-1} [-H^{0i}\theta_i + H^{ij}\theta_i \theta_j] (a\partial_a) (b\partial_b - a\partial_a) + H^{ij} \tilde{\phi}_i \tilde{\phi}_j \\ &\quad + 2a^{-1} [H^{0i} - H^{ij}\theta_j] \tilde{\phi}_i (b\partial_b - a\partial_a) + 2[-H^{ij}\theta_j] \tilde{\phi}_i (a\partial_a) \\ &\quad + a^{-2} [H^{00} - 2H^{i0}\theta_i + H^{ij}\theta_i \theta_j + aH^{ij}(-\delta_{ij} + n\theta_i \theta_j) - (n-1)aH^{0i}\theta_i] (b\partial_b - a\partial_a) \\ &\quad + [H^{ij}(-\delta_{ij} + (n+1)\theta_i \theta_j)] a\partial_a + [-nH^{ij}\theta_j] \tilde{\phi}_i, \\ &= \square_{\tilde{m}} - a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} \tilde{h}_{\rho\rho} [(b\partial_b - a\partial_a)^2 + (b\partial_b - a\partial_a)] + a^{\frac{n-3}{2}} b^{\frac{n-1}{2}} \Theta_1(\tilde{h}) (\tilde{\partial}^2 + \tilde{\partial}), \\ \gamma &= \gamma_0 + [H^{ij}(-\frac{n-1}{2}\delta_{ij} + \frac{(n-1)(n+3)}{4}\theta_i \theta_j)], \quad \gamma_0 = -\frac{(n-1)(n-3)}{4}.\end{aligned}$$

- In Ω_3 , $\rho = \frac{1}{r}$, $\tau = t - r$, $\tilde{\rho} = \rho$,

$$\begin{aligned}\square_{\tilde{m}} &= 2\partial_\rho \partial_\tau + (\rho \partial_\rho)^2 + \rho \partial_\rho + \triangle_\theta, \\ \square_{\tilde{g}} &= \square_{\tilde{m}} + \rho^{-2} [H^{00} - 2H^{i0}\theta_i + H^{ij}\theta_i \theta_j] \partial_\tau^2 + [H^{ij}\theta_i \theta_j] (\rho \partial_\rho)^2 + H^{ij} \tilde{\phi}_i \tilde{\phi}_j \\ &\quad + 2[-H^{ij}\theta_j] (\rho \partial_\rho) \tilde{\phi}_i + 2\rho^{-1} [-H^{i0}\theta_i + H^{ij}\theta_i \theta_j] \partial_\tau (\rho \partial_\rho) \\ &\quad + 2\rho^{-1} [H^{i0} - H^{ij}\theta_j] \partial_\tau \tilde{\phi}_i + [H^{ij}(-\delta_{ij} + (n+1)\theta_i \theta_j)] \rho \partial_\rho \\ &\quad + \rho^{-1} [-(n-1)H^{0i}\theta_i + H^{ij}(-\delta_{ij} + n\theta_i \theta_j)] \partial_\tau + n[-H^{ij}\theta_j] \tilde{\phi}_i \\ &= \square_{\tilde{m}} - \rho^{\frac{n-5}{2}} \tilde{h}_{\rho\rho} \partial_\tau^2 + \rho^{\frac{n-3}{2}} \Theta_1(\tilde{h}) (\tilde{\partial}^2 + \tilde{\partial}), \\ \gamma &= \gamma_0 + [H^{ij}(-\frac{n-1}{2}\delta_{ij} + \frac{(n-1)(n+3)}{4}\theta_i \theta_j)], \quad \gamma_0 = -\frac{(n-1)(n-3)}{4}.\end{aligned}$$

- In Ω_4 , $\bar{a} = 1 - \frac{1}{t}$, $\bar{b} = \frac{1}{t-r}$, $\tilde{\rho} = \bar{a}\bar{b}$,

$$\begin{aligned}
\Box_{\tilde{m}} &= -2\partial_{\bar{a}}\bar{b}\partial_{\bar{b}} + \bar{a}(2 - \bar{a})\partial_{\bar{a}}^2 + 2\partial_{\bar{a}} - (n+1)\bar{a}\partial_{\bar{a}} + \frac{1}{(1-\bar{a})^2}\Delta_{\theta} - \frac{n-1}{1-\bar{a}}(\bar{a}\partial_{\bar{a}} - \bar{b}\partial_{\bar{b}}) \\
&= 2\partial_{\bar{a}}(\bar{a}\partial_{\bar{a}} - \bar{b}\partial_{\bar{b}}) - (\bar{a}\partial_{\bar{a}})^2 - n\bar{a}\partial_{\bar{a}} + \frac{1}{(1-\bar{a})^2}\Delta_{\theta} - \frac{n-1}{1-\bar{a}}(\bar{a}\partial_{\bar{a}} - \bar{b}\partial_{\bar{b}}) \\
\Box_{\tilde{g}} &= \Box_{\tilde{m}} + \bar{a}^{-2}[H^{00} - 2H^{i0}\theta_i + H^{ij}\theta_i\theta_j](\bar{b}\partial_{\bar{b}} - \bar{a}\partial_{\bar{a}})^2 + H^{00}(\bar{a}\partial_{\bar{a}})^2 \\
&\quad + (1-\bar{a})^{-2}H^{ij}\partial_i\partial_j + 2\bar{a}^{-1}[H^{00} - H^{0i}\theta_i](\bar{b}\partial_{\bar{b}} - \bar{a}\partial_{\bar{a}}) \\
&\quad - 2(1-\bar{a})^{-1}H^{0i}(\bar{a}\partial_{\bar{a}})\partial_i + 2(\bar{a}(1-\bar{a}))^{-1}[-H^{i0} + H^{ij}\theta_j](\bar{b}\partial_{\bar{b}} - \bar{a}\partial_{\bar{a}})\partial_i \\
&\quad + (1-\bar{a})^{-2}[-H^{ij}\theta_j - (n-1)H^{0i}]\partial_i + nH^{00}\bar{a}\partial_{\bar{a}} + \bar{a}^{-2}[H^{00} - 2H^{i0}\theta_i + H^{ij}\theta_i\theta_j \\
&\quad + (n-1)\bar{a}(H^{00} - H^{0i}\theta_i) - \bar{a}(1-\bar{a})^{-1}H^{ij}(-\delta_{ij} + \theta_i\theta_j)](\bar{b}\partial_{\bar{b}} - \bar{a}\partial_{\bar{a}}) \\
&= \Box_{\tilde{m}} - \bar{a}^{\frac{n-5}{2}}\bar{b}^{\frac{n-1}{2}}\tilde{h}_{\rho\rho}[(\bar{b}\partial_{\bar{b}} - \bar{a}\partial_{\bar{a}})^2 + (\bar{b}\partial_{\bar{b}} - \bar{a}\partial_{\bar{a}})] + \bar{a}^{\frac{n-3}{2}}\bar{b}^{\frac{n-1}{2}}\Theta_1(\tilde{h})(\tilde{\partial}^2 + \tilde{\partial}), \\
\gamma &= \gamma_0 + \frac{n^2-1}{4}H^{00}, \quad \gamma_0 = -\frac{n^2-1}{4}.
\end{aligned}$$

- In Ω_5 , $\phi = \frac{1}{t}$, $y = \frac{x}{t}$, $\tilde{\rho} = \phi$,

$$\begin{aligned}
\Box_{\tilde{m}} &= -(\phi\partial_{\phi} + \sum_i y^i\partial_{y^i})^2 - n(\phi\partial_{\phi} + \sum_i y^i\partial_{y^i}) + \Delta_y, \\
\Box_{\tilde{g}} &= \Box_{\tilde{m}} + H^{00}(\phi\partial_{\phi} + y^k\partial_{y^k})^2 + H^{ij}\partial_{y^i}\partial_{y^j} - 2H^{0i}(\phi\partial_{\phi} + y^k\partial_{y^k})\partial_{y^i} \\
&\quad + nH^{00}(\phi\partial_{\phi} + y^i\partial_{y^i}) - (n+1)H^{0i}\partial_{y^i}, \\
&= \Box_{\tilde{m}} + \phi^{\frac{n-1}{2}}\Theta_1(\tilde{h})(\tilde{\partial}^2 + \tilde{\partial}), \\
\gamma &= \gamma_0 + \frac{n^2-1}{4}H^{00}, \quad \gamma_0 = -\frac{n^2-1}{4}.
\end{aligned}$$

We finish the proof. $\square$

Next, we consider the commutator of $\Box_{\tilde{m}}$ and a basis of $\mathcal{V}_b(X)$. By Lemma 2.1, we can choose a basis of $\mathcal{V}_b(\Omega_j)$, denoted by $\mathcal{B}_j$, for $0 \leq j \leq 5$ as follows:

$$(4.4) \quad \begin{cases} \mathcal{B}_0 = \{\partial_{\mu} : \mu = 0, 1, \dots, n\}, \\ \mathcal{B}_j = \{Z_{\mu\nu} : \mu, \nu = 0, 1, \dots, n, j=1, 2, 4, 5\}, \\ \mathcal{B}_3 = \{\partial_{\tau}, \rho\partial_{\rho}, Z_{ij} : i, j = 1, \dots, n\}. \end{cases}$$

Lemma 4.7. *Suppose (t, x) are harmonic coordinates w.r.t. g . Then in each domain Ω_j for $1 \leq j \leq 5$, with the choice of $\tilde{\rho}$ specified in Section 2,*

$$[\Box_{\tilde{m}}, Z_{00}] = [\Box_{\tilde{m}}, Z_{ij}] = 0, \quad [\Box_{\tilde{m}}, Z_{0i}] = c_i(\Box_{\tilde{m}} + \gamma_0) + \tilde{\partial} + c'_i,$$

for $1 \leq i \leq n$, where $c_i, c'_i \in C^\infty(\Omega_j)$ and $\tilde{\partial} \in \mathcal{V}_b(\Omega_j)$. In particular, in Ω_3 ,

$$[\Box_{\tilde{m}}, \partial_{\tau}] = 0, \quad [\Box_{\tilde{m}}, \rho\partial_{\rho}] = \Box_{\tilde{m}} - (\rho\partial_{\rho})^2 - \rho\partial_{\rho} - \Delta_{\theta}.$$

Proof. For any $u \in C^\infty(X)$,

$$\begin{aligned}
[\Box_{\tilde{m}}, Z_{\mu\nu}]u &= (\Box_{\tilde{m}} + \gamma_0)Z_{\mu\nu}u - Z_{\mu\nu}(\Box_{\tilde{m}} + \gamma_0)u \\
&= \tilde{\rho}^{-\frac{n+3}{2}}\Box_{\tilde{m}}\tilde{\rho}^{\frac{n-1}{2}}Z_{\mu\nu}u - Z_{\mu\nu}\tilde{\rho}^{-\frac{n+3}{2}}\Box_{\tilde{m}}\tilde{\rho}^{\frac{n-1}{2}}u \\
&= \tilde{\rho}^{-\frac{n+3}{2}}[\Box_{\tilde{m}}, Z_{\mu\nu}]\tilde{\rho}^{\frac{n-1}{2}}u + \tilde{\rho}^{-\frac{n+3}{2}}\Box_{\tilde{m}}[\tilde{\rho}^{\frac{n-1}{2}}, Z_{\mu\nu}]u - [Z_{\mu\nu}, \tilde{\rho}^{-\frac{n+3}{2}}]\Box_{\tilde{m}}\tilde{\rho}^{\frac{n-1}{2}}u \\
&= c_{\mu\nu}(\Box_{\tilde{m}} + \gamma_0)u - \frac{n-1}{2}(\Box_{\tilde{m}} + \gamma_0)\tilde{\rho}^{-1}Z_{\mu\nu}(\tilde{\rho})u + \frac{n+3}{2}\tilde{\rho}^{-1}Z_{\mu\nu}(\tilde{\rho})(\Box_{\tilde{m}} + \gamma_0)u \\
&= (c_{\mu\nu} + 2\tilde{\rho}^{-1}Z_{\mu\nu}(\tilde{\rho}))(\Box_{\tilde{m}} + \gamma_0)u - \frac{n-1}{2}[\Box_{\tilde{m}}, \tilde{\rho}^{-1}Z_{\mu\nu}(\tilde{\rho})]u
\end{aligned}$$

Recall that $[\Box_m, Z_{\mu\nu}] = c_{\mu\nu}\Box_m$ with $c_{0i} = c_{ij} = 0$, $c_{00} = 2$ and

$$\rho^{-1}Z_{00}(\rho) = \phi^{-1}Z_{00}(\phi) = -1, \quad \rho^{-1}Z_{ij}(\rho) = \phi^{-1}Z_{ij}(\phi) = 0.$$

Hence $[\square_{\tilde{m}}, Z_{00}] = [\square_{\tilde{m}}, Z_{ij}] = 0$. Notice that we choose $\tilde{\rho} = \rho$ in $\Omega_1, \Omega_2, \Omega_3$ and $\tilde{\rho} = \phi$ in Ω_4, Ω_5 and

$$\rho^{-1}Z_{0i}(\rho) = -\frac{t\theta_i}{r} = \begin{cases} -s\theta_i & \text{in } \Omega_1 \\ -(1-a)\theta_i & \text{in } \Omega_2, \\ -(1+\rho\tau)\theta_i & \text{in } \Omega_3 \end{cases}, \quad \phi^{-1}Z_{00}(\phi) = -\frac{x^i}{t} = \begin{cases} -(1-\bar{a})\theta_i & \text{in } \Omega_4 \\ -y^i & \text{in } \Omega_5 \end{cases},$$

Hence $\tilde{\rho}^{-1}Z_{0i}(\tilde{\rho}) \in C^\infty(\Omega_j)$ for all $1 \leq i \leq n$ and $1 \leq j \leq 5$. Since

$$[\mathcal{V}_b(\Omega_j), C^\infty(\Omega_j)] \subset C^\infty(\Omega_j)$$

It is obviously for $j = 1$ or $j = 5$,

$$[\square_{\tilde{m}}, \tilde{\rho}^{-1}Z_{0i}(\tilde{\rho})] \in \mathcal{V}_b(\Omega_j) + C^\infty(\Omega_j).$$

For $j = 2, 3, 4$, by Lemma 4.6, we only need to consider

$$[2\partial_{\rho_1}D, \tilde{\rho}^{-1}Z_{0i}(\tilde{\rho})] = 2\partial_{\rho_1}D(\tilde{\rho}^{-1}Z_{0i}(\tilde{\rho})) + 2D(\tilde{\rho}^{-1}Z_{0i}(\tilde{\rho}))\partial_{\rho_1} + 2\partial_{\rho_1}(\tilde{\rho}^{-1}Z_{0i}(\tilde{\rho}))D$$

where $2\partial_{\rho_1}D(\tilde{\rho}^{-1}Z_{0i}(\tilde{\rho})), 2\partial_{\rho_1}(\tilde{\rho}^{-1}Z_{0i}(\tilde{\rho})) \in C^\infty(\Omega_j)$ and $D(\tilde{\rho}^{-1}Z_{0i}(\tilde{\rho})) \in \rho_1 C^\infty(\Omega_j)$. Hence

$$[2\partial_{\rho_1}D, \tilde{\rho}^{-1}Z_{0i}(\tilde{\rho})] \in \mathcal{V}_b(\Omega_j) + C^\infty(\Omega_j).$$

We finish the proof. $\square$

4.4. Conformal Transformation of Reduced Einstein Equations. With the conformal transformation (4.1), the reduced Einstein vacuum equations (1.3) is equivalent to the following system:

$$(4.5) \quad (\square_{\tilde{g}} + \gamma)\tilde{h}_{\mu\nu} = (\rho_0\rho_2)^{\frac{n-1}{2}}\rho_1^{\frac{n-5}{2}}\tilde{F}_{\mu\nu}(\tilde{h}, \tilde{\partial}\tilde{h}),$$

where $\gamma = \gamma_0 + \tilde{\rho}^{\frac{n-1}{2}}\Theta_1(\tilde{h})$ is given in Lemma 4.6 and

$$\tilde{F}_{\mu\nu} = (\rho_0\rho_2)^{-(n+1)}\rho_1^{-(n-1)}F_{\mu\nu}(\tilde{\rho}^{\frac{n-1}{2}}\tilde{h})(\partial(\tilde{\rho}^{\frac{n-1}{2}}\tilde{h}), \partial(\tilde{\rho}^{\frac{n-1}{2}}\tilde{h})).$$

Using the explicit formula of $F_{\mu\nu}$ given in [LR1], we have

Lemma 4.8. For $n \geq 3$,

$$\tilde{F}_{\mu\nu} = \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h}, \tilde{\partial}\tilde{h}) + \rho_1\Theta_1(\tilde{h})(\tilde{\partial}\tilde{h}) + \rho_1\Theta_2(\tilde{h}).$$

In particular, in domains $\Omega_2, \Omega_3, \Omega_4$,

$$\tilde{F}_{\mu\nu} = \theta_\mu\theta_\nu(\frac{1}{4}(D\text{tr}_m\tilde{h})^2 - \frac{1}{2}D\tilde{h}_\beta^\alpha D\tilde{h}_\alpha^\beta) + \rho_1(\Theta_0(\tilde{h})(\tilde{h})(\tilde{\partial}\tilde{h}, \tilde{\partial}\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})) + \rho_1^2\Theta_2(\tilde{h})$$

where $D \in \mathcal{V}_b(\Omega_i)$ is defined in Lemma 4.3.

Proof. By [LR1], $F_{\mu\nu}(\partial h, \partial h) = P_{\mu\nu}(\partial h, \partial h) + Q_{\mu\nu}(\partial h, \partial h) + G(h)_{\mu\nu}(\partial h, \partial h)$, where

$$P_{\mu\nu}(\partial h, \partial h) = \frac{1}{4}\partial_\mu(\text{tr}_m h)\partial_\nu(\text{tr}_m h) - \frac{1}{2}\partial_\mu h_\alpha^\beta \partial_\nu h_\beta^\alpha,$$

$Q_{\mu\nu}(\partial h, \partial h)$ satisfies the null condition and

$$G(h)_{\mu\nu}(\partial h, \partial h) = \Theta_1(h)(\partial h, \partial h)$$

First notice that $[\partial, \tilde{\rho}^{\frac{n+1}{2}}] \in \tilde{\rho}^{\frac{n+1}{2}}\mathcal{A}^{0,0,0}(X)$. Hence by the conformal transformation (4.1),

$$P_{\mu\nu}(\partial h, \partial h) = \tilde{\rho}^{n-1}((\rho_0\rho_2)^2\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h}, \tilde{\partial}\tilde{h}) + \rho_0\rho_2\tilde{\rho}\Theta_1(\tilde{h})(\tilde{\partial}\tilde{h}) + \tilde{\rho}^2\Theta_2(\tilde{h})),$$

$$Q_{\mu\nu}(\partial h, \partial h) = \tilde{\rho}^n\rho_0\rho_2(\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h}, \tilde{\partial}\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})) + \tilde{\rho}^{n+1}\Theta_2(\tilde{h}),$$

$$G(h)_{\mu\nu}(\partial h, \partial h) = \tilde{\rho}^{\frac{3n-3}{2}}((\rho_0\rho_2)^2\Theta_1(\tilde{h})(\tilde{\partial}\tilde{h}, \tilde{\partial}\tilde{h}) + \rho_0\rho_2\tilde{\rho}\Theta_2(\tilde{h})(\tilde{\partial}\tilde{h}) + \tilde{\rho}^2\Theta_3(\tilde{h})).$$

In particular, in domain $\Omega_2, \Omega_3, \Omega_4$,

$$\begin{aligned} P_{\mu\nu}(\partial h, \partial h) &= \tilde{\rho}^{n-1}(\rho_0\rho_2)^2\theta_\mu\theta_\nu(\frac{1}{4}(D\text{tr}_m\tilde{h})^2 - \frac{1}{2}D\tilde{h}_\beta^\alpha D\tilde{h}_\alpha^\beta) \\ &\quad + \tilde{\rho}^n\rho_0\rho_2(\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h}, \tilde{\partial}\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})) + \tilde{\rho}^{n+1}\Theta_2(\tilde{h}) \end{aligned}$$

We finish the proof. $\square$

Lemma 4.9. *The equations (4.5) implies that for all $k \geq 0$*

$$(4.6) \quad \begin{aligned} & (\square_{\tilde{g}} + \gamma_0) \tilde{\partial}^k \tilde{h}_{\mu\nu} = [\square_{\tilde{m}}, \tilde{\partial}^k] \tilde{h}_{\mu\nu} + \tilde{f}_{\mu\nu}^k(\tilde{h}), \quad \text{where} \\ & \tilde{f}_{\mu\nu}^k(h) = (\rho_0 \rho_2)^{\frac{n-1}{2}} \rho_1^{\frac{n-5}{2}} \sum_{\alpha_1 + \dots + \alpha_l \leq k+2, \ 2 \leq l \leq k+2, \ 0 \leq \alpha_i \leq k+1} \Theta_0(\tilde{h})(\tilde{\partial}^{\alpha_1} \tilde{h}, \dots, \tilde{\partial}^{\alpha_l} \tilde{h}). \end{aligned}$$

Proof. For $k = 0$, obviously

$$\begin{aligned} \tilde{f}_{\mu\nu}^0(\tilde{h}) &= (\gamma - \gamma_0) \tilde{h}_{\mu\nu} + (\rho_0 \rho_2)^{\frac{n-1}{2}} \rho_1^{\frac{n-5}{2}} \tilde{F}_{\mu\nu} \\ &= (\rho_0 \rho_2)^{\frac{n-1}{2}} \rho_1^{\frac{n-5}{2}} \sum_{\alpha_1 + \alpha_2 \leq 2, 0 \leq \alpha_i \leq 1} \Theta_0(\tilde{h})(\tilde{\partial}^{\alpha_1} \tilde{h}, \tilde{\partial}^{\alpha_2} \tilde{h}). \end{aligned}$$

For $k \geq 1$, $(\square_{\tilde{g}} + \gamma_0) \tilde{\partial}^k \tilde{h}_{\mu\nu} = \tilde{\partial}^k \tilde{f}_{\mu\nu}^0(\tilde{h}) + [\square_{\tilde{g}}, \tilde{\partial}^k] \tilde{h}_{\mu\nu}$. Hence

$$\tilde{f}_{\mu\nu}^k(\tilde{h}) = [\square_{\tilde{g}} - \square_{\tilde{m}}, \tilde{\partial}^k] \tilde{h}_{\mu\nu} + \tilde{\partial}^k \tilde{f}_{\mu\nu}^0(\tilde{h})$$

By Lemma 4.6, $\tilde{f}_{\mu\nu}^k(\tilde{h})$ has the form in (4.6). $\square$

Lemma 4.10. *Let $\tilde{D}$ be any vector field in $\mathcal{B}_j$ for $1 \leq j \leq 5$ and I a multi-index. Then in each domain Ω_j for $1 \leq j \leq 5$, the equations (4.6) implies that*

$$(4.7) \quad \begin{aligned} & (\square_{\tilde{g}} + \gamma_0) \tilde{D}^I \tilde{h}_{\mu\nu} = \sum_{0 \leq i \leq |I|+1} \tilde{\partial}^i \tilde{h}_{\mu\nu} + \tilde{f}_{\mu\nu}^I(\tilde{h}), \quad \text{where} \\ & \tilde{f}_{\mu\nu}^I(\tilde{h}) = (\rho_0 \rho_2)^{\frac{n-1}{2}} \rho_1^{\frac{n-5}{2}} \sum_{\alpha_1 + \dots + \alpha_l \leq |I|+2, \ 2 \leq l \leq |I|+2, \ 0 \leq \alpha_i \leq |I|+1} \Theta_0(\tilde{h})(\tilde{\partial}^{\alpha_1} \tilde{h}, \dots, \tilde{\partial}^{\alpha_l} \tilde{h}). \end{aligned}$$

Proof. It is obviously true if $|I| = 0$ by Lemma 4.9. Then for $|I| > 0$, by induction,

$$\begin{aligned} [\square_{\tilde{m}}, \tilde{D}^I] \tilde{h}_{\mu\nu} &= \sum_{|K|+|J|=|I|-1} C_{K,J} \tilde{D}^K [\square_{\tilde{m}}, \tilde{D}] \tilde{D}^J \tilde{h}_{\mu\nu} \\ &= \sum_{|K|+|J|=|I|-1} C_{K,J} \tilde{D}^K (c(\square_{\tilde{g}} + \gamma_0) + c(\square_{\tilde{m}} - \square_{\tilde{g}}) + \tilde{\partial}^2 + \tilde{\partial} + c') \tilde{D}^J \tilde{h}_{\mu\nu} \\ &= \sum_{|K|+|J|=|I|-1} C_{K,J} \tilde{D}^K \left(\sum_{0 \leq i \leq |J|+2} \tilde{\partial}^i \tilde{h}_{\mu\nu} + c \tilde{f}_{\mu\nu}^J(\tilde{h}) + (\rho_0 \rho_2)^{\frac{n-1}{2}} \rho_1^{\frac{n-5}{2}} \Theta_0(\tilde{h})(\tilde{\partial}^2 + \tilde{\partial}) \tilde{D}^J \tilde{h}_{\mu\nu} \right) \end{aligned}$$

where $C_{K,J}$ are constants, $c, c' \in C^\infty(\Omega_j)$. Hence $[\square_{\tilde{m}}, \tilde{D}^I] \tilde{h}_{\mu\nu}$ can be expressed as

$$\sum_{0 \leq i \leq |I|+1} \tilde{\partial}^i \tilde{h}_{\mu\nu} + (\rho_0 \rho_2)^{\frac{n-1}{2}} \rho_1^{\frac{n-5}{2}} \sum_{\alpha_1 + \dots + \alpha_l \leq |I|+2, \ 2 \leq l \leq |I|+2, \ 0 \leq \alpha_i \leq |I|+1} \Theta_0(\tilde{h})(\tilde{\partial}^{\alpha_1} \tilde{h}, \dots, \tilde{\partial}^{\alpha_l} \tilde{h}).$$

we prove the lemma. $\square$

5. TIME-LIKE FUNCTIONS.

Time-like function is the most important concept associated to a Lorentzian metric: we use it to define the positive quadratic form on space-like hypersurfaces, as well as the energy norms. In this section, we consider the time-like functions w.r.t. $\tilde{m} = \tilde{\rho}^2 m$ and $\tilde{g} = \tilde{\rho}^2 g = \tilde{m} + \tilde{\rho}^{\frac{n+3}{2}} \tilde{h}$ with $\tilde{h}$ small on the interior of X , denote by $\tilde{X}$.

Definition 4. *Suppose $\tilde{g}$ extends to a Lorentz b -metric on X^2 of signature $(n, 1)$, $T \in C^1(\tilde{X})$ and $p \in \tilde{X}$. We say T is time-like w.r.t. $\tilde{g}$ at p iff*

$$\langle \nabla T, \nabla T \rangle_{\tilde{g}} < 0 \quad \text{at } p,$$

and null w.r.t. $\tilde{g}$ at p iff

$$\langle \nabla T, \nabla T \rangle_{\tilde{g}} = 0 \quad \text{at } p.$$

A hypersurface $\Sigma \subset X$ is called space-like (resp. null) w.r.t. $\tilde{g}$ iff Σ has a defining function $T \in C^1(\tilde{X})$ such that T is time-like (resp. null) on Σ .

Definition 5. A quadratic form field associated to the time-like function T w.r.t. $\tilde{g}$ is defined by

$$\mathcal{F}_{\tilde{g}}(T, v) = \langle \nabla T, \nabla v \rangle_{\tilde{g}} \nabla v - \frac{1}{2} \langle \nabla v, \nabla v \rangle_{\tilde{g}} \nabla T + \frac{1}{2} \gamma_0 v^2 \nabla T.$$

Here ∇ is the covariant derivative w.r.t. $\tilde{g}$ and $\gamma_0 < 0$ is some constant.

Lemma 5.1. Suppose T, T' are two time-like functions w.r.t. a Lorentzian metric $\tilde{g}$ at $p \in \mathring{X}$ such that $\langle \nabla T, \nabla T' \rangle_{\tilde{g}} < 0$ at p , then $\langle \mathcal{F}_{\tilde{g}}(T, v), \nabla T' \rangle_{\tilde{g}}$ is a strict positive quadratic form of $(\nabla v, v)$ at p .

See Chapter 6 in [Ho1] for proof in details. Moreover, we have the following formulas from the divergence theorem directly, which implies the energy estimates if choosing T and Ω properly.

Lemma 5.2. Suppose that the domain $\Omega \subset \mathring{X}$ has piecewise smooth boundary $\partial\Omega$ with defining function T' and T is a C^2 time-like function w.r.t. $\tilde{g}$ on $\bar{\Omega} \cap \mathring{X}$. Assume $T'' \in C^1(\bar{\Omega} \cap \mathring{X})$. Then

$$\int_{\Omega} \operatorname{div}_{\tilde{g}}(e^{-2T''} \mathcal{F}_{\tilde{g}}(T, v)) d\operatorname{vol}_{\tilde{g}} = \int_{\partial\Omega} e^{-2T''} \langle \mathcal{F}_{\tilde{g}}(T, v), \nabla T' \rangle_{\tilde{g}} d\mu_{\tilde{g}}^{T'}.$$

Here $d\mu_{\tilde{g}}^{T'}$ is the volume form on $\partial\Omega$ such that $dT' \wedge d\mu_{\tilde{g}}^{T'} = d\operatorname{vol}_{\tilde{g}}$ and $\partial\Omega$ is oriented by $d\mu_{\tilde{g}}^{T'} > 0$, and ∇ is the covariant derivative w.r.t. $\tilde{g}$ and

$$(5.1) \quad \begin{aligned} \operatorname{div}_{\tilde{g}}(e^{-2T''} \mathcal{F}_{\tilde{g}}(T, v)) &= e^{-2T''} \left(-2 \langle \mathcal{F}_{\tilde{g}}(T, v), \nabla T'' \rangle_{\tilde{g}} + \langle \nabla v, \nabla T \rangle_{\tilde{g}} (\square_{\tilde{g}} + \gamma_0) v + \mathcal{Q}_{\tilde{g}}(T, v) \right), \\ \text{where } \mathcal{Q}_{\tilde{g}}(T, v) &= \frac{1}{2} (\gamma_0 v^2 - \langle \nabla v, \nabla v \rangle_{\tilde{g}}) \square_{\tilde{g}} T + \nabla^2 T (dv, dv). \end{aligned}$$




FIGURE 3. Division of $X_{t \geq 0}$ by space like hypersurfaces and null hypersurfaces.

For the purpose of energy estimates, we divide $X_{t \geq 0}$ into four domains $\{\tilde{\Omega}_i\}_{1 \leq i \leq 4}$ by space-like hypersurfaces and null hypersurface w.r.t. $\tilde{m}$, or equivalently w.r.t. $\tilde{g}$. See Figure 3. We list the time-like functions used to define those surfaces, which are also used to define the energy norms in next section, domain by domain. For $1 \leq i \leq 4$, we use T_i, T'_i to denote time-like functions in $\tilde{\Omega}_i \cap \mathring{X}$ such that T'_i extends to a null functions on $S_1^+ \cap \tilde{\Omega}_i$. We use T''_i to denote the weight functions if necessary. We also compute the corresponding quadratic form field $\mathcal{F}_{\tilde{g}}(T_i, v)$, quadratic forms $\langle \mathcal{F}_{\tilde{g}}(T_i, v), \nabla T'_i \rangle_{\tilde{g}}$, $\langle \mathcal{F}_{\tilde{g}}(T_i, v), \nabla T''_i \rangle_{\tilde{g}}$ and $\mathcal{Q}_{\tilde{g}}(T_i, v)$ for $1 \leq i \leq 4$.

Through out this section, we assume the following:

- (A1) (t, x) are harmonic coordinates w.r.t. $g = m + \tilde{\rho}^{\frac{n-1}{2}} \tilde{h}$.
- (A2) $|(\rho_0 \rho_2)^{\frac{n-1}{2}} \tilde{h}|$, $|(\rho_0 \rho_2)^{\frac{n-1}{2}} \tilde{\partial} \tilde{h}|$, $|(\rho_0 \rho_2)^{\frac{n-1}{2}} \rho_1^{-\delta} \tilde{h}_{\rho\rho}|$, $|(\rho_0 \rho_2)^{\frac{n-1}{2}} \rho_1^{-\delta} \tilde{\partial} \tilde{h}_{\rho\rho}|$ are small enough for some $0 < \delta < \frac{1}{2}$.
- (A3) $\delta' \in (0, \delta)$ satisfies $\frac{1}{2} - \delta' < 1 - 2\delta$ and $\alpha > 1$.

5.1. **In $\tilde{\Omega}_1$.** In $\Omega_1 \cup \Omega_0$, define

$$T_1 = \frac{t}{\psi_1(r)};$$

$$\tilde{\Omega}_1 = \{0 \leq T_1 \leq \frac{3}{4}\}, \quad \Sigma_0 = \{T_1 = 0\}, \quad \Sigma_1 = \{T_1 = \frac{3}{4}\}.$$

where $\psi_1 \in C^2(\tilde{\Omega}_1)$ is defined by

$$\psi_1(r) = \begin{cases} \frac{3}{8} + \frac{3}{4}r^2 - \frac{1}{8}r^4 & \text{for } r \leq 1 \\ r & \text{for } r > 1 \end{cases}$$

Hence $0 \leq \psi_1' \leq 1$. Then in $\tilde{\Omega}_1$,

$$\langle \nabla T_1, \nabla T_1 \rangle_{\tilde{m}} = (\tilde{\rho}\psi_1(r))^{-2}(-1 + T^2(\psi_1'(r))^2) \leq -\frac{7}{16}(\tilde{\rho}\psi_1(r))^{-2} < -C$$

for some constant $C > 0$. Hence T_1 is a regular time-like function w.r.t. $\tilde{m}$ all over $\tilde{\Omega}_1$ and bounded. We choose $\tilde{\rho} = \rho_0 = \psi_1^{-1}$ and $\rho_1 = \rho_2 = 1$ in $\tilde{\Omega}_1$. Here $\tilde{\Omega}_1 \cap \Omega_0$ is bounded in the interior of X . Near the boundary of $\tilde{\Omega}_1$, i.e. in $\tilde{\Omega}_1 \cap \Omega_1$, recall that we use the following coordinates and boundary defining functions:

$$(s, \rho, \theta) = (\frac{t}{r}, \frac{1}{r}, \frac{x}{r}); \quad \tilde{\rho} = \rho_0 = \rho, \quad \rho_1 = \rho_2 = 1.$$

Let $\gamma_0 < 0$ be a constant and we have the following two lemmas.

Lemma 5.3. In $\tilde{\Omega}_1 \cap \Omega_1$,

$$\langle \mathcal{F}_{\tilde{m}}(T_1, v), \nabla T_1 \rangle_{\tilde{m}} = \frac{1}{2}|(1-s^2)\partial_s v - \rho\partial_\rho v|^2 + \frac{1}{2}|\rho\partial_\rho v|^2 + \frac{1}{2}(1-s^2)(|\partial_\theta v|^2 - \gamma_0 v^2),$$

$$\mathcal{Q}_{\tilde{m}}(T_1, v) = -s(|\partial_\theta v|^2 - \gamma_0 v^2).$$

Here ∇ is the covariant derivative w.r.t. $\tilde{m}$.

Proof. In $\tilde{\Omega}_1 \cap \Omega_1$, change variable to $\xi = -\ln \rho \in (0, \infty)$ and hence $\frac{d\rho}{\rho} = -d\xi$ and $\rho\partial_\rho = -\partial_\xi$. Then in coordinates $\{s, \xi, \theta\}$,

$$\tilde{m} = -ds^2 - 2sdsd\xi + (1-s^2)d^2\xi + d^2\theta.$$

It is a product type metric. WLOG, at any fixed point $p = (s, \xi, \theta)$, let θ denote the normal coordinate w.r.t. standard spherical metric $d^2\theta$ on $\mathbb{S}^{n-1}$. Then at p , we can write the metric components as follows:

$$\tilde{m} = \begin{bmatrix} -1 & -s & 0 \\ -s & 1-s^2 & 0 \\ 0 & 0 & \mathbb{1}_{n-1} \end{bmatrix}, \quad \tilde{m}^{-1} = \begin{bmatrix} -1+s^2 & -s & 0 \\ -s & 1 & 0 \\ 0 & 0 & \mathbb{1}_{n-1} \end{bmatrix}.$$

For the quadratic form, notice that

$$\langle \mathcal{F}_{\tilde{m}}(T_1, v), \nabla T_1 \rangle_{\tilde{m}} = \langle \nabla T_1, \nabla v \rangle_{\tilde{m}}^2 - \frac{1}{2}\langle \nabla v, \nabla v \rangle_{\tilde{m}} \langle \nabla T_1, \nabla T_1 \rangle_{\tilde{m}} + \frac{1}{2}\gamma_0 \langle \nabla T_1, \nabla T_1 \rangle_{\tilde{m}} v^2,$$

where

$$\langle \nabla T_1, \nabla T_1 \rangle_{\tilde{m}} = -1 + s^2,$$

$$\langle \nabla T_1, \nabla v \rangle_{\tilde{m}} = (-1 + s^2)\partial_s v - s\partial_\xi v,$$

$$\langle \nabla v, \nabla v \rangle_{\tilde{m}} = (-1 + s^2)|\partial_s v|^2 - 2s\partial_s v \partial_\xi v + |\partial_\xi v|^2 + |\partial_\theta v|^2.$$

Then the formula for $\langle \mathcal{F}_{\tilde{m}}(T_1, v), \nabla T_1 \rangle_{\tilde{m}}$ follows directly. For $\mathcal{Q}_{\tilde{m}}(T_1, v)$, by Lemma 5.2 we only need to compute $\square_{\tilde{m}} T_1$ and $\nabla^2 T_1(dv, dv)$. Using to the formula of $\square_{\tilde{m}}$ in the proof of Lemma 4.6,

$$\square_{\tilde{m}} T_1 = 2s; \quad \text{and}$$

$$\nabla^2 T_1(dv, dv) = (\partial_I \partial_J T_1 - \Gamma_{IJ}^K \partial_K T_1) \nabla^I v \nabla^J v.$$

Here $I, J, K \in \{s, \xi, \theta\}$. At p , the non-zero connection components are

$$\Gamma_{ss}^s(\tilde{m}) = s, \quad \Gamma_{s\xi}^s(\tilde{m}) = s^2, \quad \Gamma_{\xi\xi}^s(\tilde{m}) = -s(1-s^2),$$

$$\Gamma_{ss}^\xi(\tilde{m}) = -1, \quad \Gamma_{s\xi}^\xi(\tilde{m}) = -s, \quad \Gamma_{\xi\xi}^\xi(\tilde{m}) = -s^2.$$

Hence

$$\nabla^2 T_1(dv, dv) = -s(1-s^2)|\partial_s v|^2 - 2s^2\partial_s v \partial_\xi v + s|\partial_\xi v|^2$$

and $\mathcal{Q}_{\tilde{m}}(T_1, v)$ follows. $\square$

Those two formulae do not change much when we take the background metric $\tilde{g}$ in the following sense:

Lemma 5.4. *In $\tilde{\Omega}_1$,*

$$\begin{aligned} \langle \mathcal{F}_{\tilde{g}}(T_1, v), \nabla T_1 \rangle_{\tilde{g}} - \langle \mathcal{F}_{\tilde{m}}(T_1, v), \nabla T_1 \rangle_{\tilde{m}} &= \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h})(v, \tilde{\partial}v), \\ \mathcal{Q}_{\tilde{g}}(T_1, v) - \mathcal{Q}_{\tilde{m}}(T_1, v) &= \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h})(v, \tilde{\partial}v) + \rho_0^{\frac{n-1}{2}} \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})(\tilde{\partial}v, \tilde{\partial}v). \end{aligned}$$

Proof. We only need to concern the region near boundary. In $\tilde{\Omega}_1 \cap \Omega_1$, let us do the similar computation as in the proof of Lemma 5.3 by using the same coordinate (s, ρ, θ) . First, by the proof of Proposition 4.1, write

$$\tilde{g} = \tilde{m} + \Theta_1(h) = \tilde{m} + \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h}), \quad \tilde{g}^{-1} = \tilde{m}^{-1} + \Theta_1(h) = \tilde{m}^{-1} + \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h}).$$

Hence the first formula follows directly by

$$\begin{aligned} \langle \nabla T_1, \nabla T_1 \rangle_{\tilde{g}} - \langle \nabla T_1, \nabla T_1 \rangle_{\tilde{m}} &= \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h}), \\ \langle \nabla T_1, \nabla v \rangle_{\tilde{g}} - \langle \nabla T_1, \nabla v \rangle_{\tilde{m}} &= \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h})(\tilde{\partial}v), \\ \langle \nabla v, \nabla v \rangle_{\tilde{g}} - \langle \nabla v, \nabla v \rangle_{\tilde{m}} &= \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h})(\tilde{\partial}v, \tilde{\partial}v). \end{aligned}$$

According to the formula of $\square_{\tilde{g}}$ in the proof of Lemma 4.6,

$$\begin{aligned} \square_{\tilde{g}} T_1 - \square_{\tilde{m}} T_1 &= \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h})(\tilde{\partial}^2 + \tilde{\partial})T_1 = \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h}); \\ \Gamma_{IJ}^K(\tilde{g}) &= \Gamma_{IJ}^K(\tilde{m}) + \rho_0^{\frac{n-1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\ \tilde{g}^{IJ} \partial_I v &= \tilde{m}^{IJ} \partial_I v + \rho_0^{\frac{n-1}{2}} \Theta_1(\tilde{h})(\tilde{\partial}v), \end{aligned}$$

which implies the second formula. Here $I, J, K \in \{s, \xi, \theta\}$. $\square$

Remark 1. *Here Lemma 5.4 also holds if (t, x) is not wave coordinates w.r.t. $\tilde{g}$.*

5.2. In $\tilde{\Omega}_2$. In Ω_2 , recall that the coordinates and boundary defining functions are as follows

$$(a, b, \theta) = (1 - \frac{t}{r}, \frac{1}{r-t}, \frac{x}{r}); \quad \rho_0 = b, \quad \rho_1 = a, \quad \rho_2 = 1, \quad \tilde{\rho} = ab.$$

Define for $\tau_0 > 8$

$$\begin{aligned} T_2 &= -\frac{2}{1+2\delta'} a^{\delta'+\frac{1}{2}} + \log b, \quad T_2' = -a, \quad T_2'' = \log b; \\ \tilde{\Omega}_2 &= \{-\frac{1}{4} \leq T_2' \leq 0, -\infty < T_2 \leq -\ln \tau_0\}, \quad \Sigma_2 = \{T_2 = -\ln \tau_0\} \cap \tilde{\Omega}_2. \end{aligned}$$

Here T_2' is a natural extension of $T_1 - 1$ into Ω_2 . Then in $\tilde{\Omega}_2$,

$$\begin{aligned} \langle \nabla T_2, \nabla T_2 \rangle_{\tilde{m}} &= -a^{\delta'-\frac{1}{2}} (2 + a^{\delta'+\frac{1}{2}} (2-a)), \\ \langle \nabla T_2', \nabla T_2' \rangle_{\tilde{m}} &= -a(2-a). \end{aligned}$$

Hence T_2, T_2' are a time-like function all over $\tilde{\Omega}_2 \cap \hat{X}$ and T_2' is null on $\tilde{\Omega}_2 \cap S_1$ w.r.t. $\tilde{m}$. In $\tilde{\Omega}_2$, with $\gamma_0 < 0$ a constant, we have the following two lemmas.

Lemma 5.5. *In $\tilde{\Omega}_2$,*

$$\begin{aligned} \langle \mathcal{F}_{\tilde{m}}(T_2, v), \nabla T_2' \rangle_{\tilde{m}} &= \frac{1}{2} a^{\delta'-\frac{1}{2}} (|b\partial_b v|^2 + |b\partial_b v - a(2-a)\partial_a v|^2) + \frac{1}{2} a(2-a) |\partial_a v|^2 \\ &\quad + \frac{1}{2} (1 + a^{\delta'+\frac{1}{2}} (2-a)) (|\partial_\theta v|^2 - \gamma_0 v^2), \\ \langle \mathcal{F}_{\tilde{m}}(T_2, v), \nabla T_2'' \rangle_{\tilde{m}} &= |\partial_a v|^2 + \frac{1}{2} a^{\delta'-\frac{1}{2}} (a(2-a) |\partial_a v|^2 + |\partial_\theta v|^2 - \gamma_0 v^2), \\ \mathcal{Q}_{\tilde{m}}(T_2, v) &= \frac{1}{2} (\frac{1}{2} - \delta') a^{\delta'-\frac{3}{2}} (|b\partial_b v|^2 + |b\partial_b v - a(2-a)\partial_a v|^2) + (1-a) |\partial_a v|^2 \\ &\quad - \frac{1}{2} a^{\delta'-\frac{1}{2}} (1 + 2\delta' - (\frac{3}{2} + \delta')a) (|\partial_\theta v|^2 - \gamma_0 v^2). \end{aligned}$$

Proof. In $\tilde{\Omega}_2$, change variable to $\xi = -\ln b \in (\ln \tau_0, \infty)$ and hence $\frac{db}{b} = -d\xi$, $b\partial_b = -\partial_\xi$. Then in coordinate (a, b, θ) ,

$$\tilde{m} = -2dad\xi + a(2-a)d^2\xi + d^2\theta.$$

With θ the normal coordinates w.r.t. standard spherical metric $d^2\theta$ at $p \in \tilde{\Omega}_2 \cap \text{Int}(X^2)$, we can write the metric components at p as follows:

$$\tilde{m} = \begin{bmatrix} 0 & -1 & 0 \\ -1 & a(2-a) & 0 \\ 0 & 0 & \mathbb{1}_{n-1} \end{bmatrix}, \quad \tilde{m}^{-1} = \begin{bmatrix} -a(2-a) & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & \mathbb{1}_{n-1} \end{bmatrix}.$$

Then the quadratic forms follow from the following computation:

$$\begin{aligned} \langle \nabla T_2, \nabla T_2' \rangle_{\tilde{m}} &= -1 - a^{\delta' + \frac{1}{2}}(2-a), \\ \langle \nabla T_2, \nabla T_2'' \rangle_{\tilde{m}} &= -a^{\delta' - \frac{1}{2}}, \\ \langle \nabla T_2, \nabla v \rangle_{\tilde{m}} &= a^{\delta' - \frac{1}{2}}(a(2-a)\partial_a v + \partial_\xi v) + \partial_a v, \\ \langle \nabla T_2', \nabla v \rangle_{\tilde{m}} &= a(2-a)\partial_a v + \partial_\xi v, \\ \langle \nabla T_2'', \nabla v \rangle_{\tilde{m}} &= \partial_a v, \\ \langle \nabla v, \nabla v \rangle_{\tilde{m}} &= -2\partial_a v \partial_\xi v - a(2-a)|\partial_a v|^2 + |\partial_\theta v|^2. \end{aligned}$$

For $\mathcal{Q}_{\tilde{m}}(T_2, v)$, we have

$$\square_{\tilde{m}} T_2 = a^{\delta' - \frac{1}{2}}(1 + 2\delta' - a(\frac{3}{2} + \delta'))$$

and the non-zero connection components at p are

$$\Gamma_{a\xi}^a(\tilde{m}) = -(1-a), \quad \Gamma_{\xi\xi}^a(\tilde{m}) = a(1-a)(2-a), \quad \Gamma_{\xi\xi}^\xi(\tilde{m}) = 1-a,$$

which implies that

$$\nabla^2 T_2(dv, dv) = (\frac{1}{2} - \delta')a^{\delta' - \frac{3}{2}}(a(2-a)\partial_a v + \partial_\xi v)^2 + (1-a)[(\partial_a v)^2 - a^{\delta' - \frac{1}{2}}(2\partial_a v \partial_\xi v + a(2-a)(\partial_a v)^2)].$$

Then $\mathcal{Q}_{\tilde{m}}(T_2, v)$ follows. $\square$

Lemma 5.6. For $n \geq 4$ and in $\tilde{\Omega}_2$,

$$\begin{aligned} \langle \mathcal{F}_{\tilde{g}}(T_2, v), \nabla T_2' \rangle_{\tilde{g}} - \langle \mathcal{F}_{\tilde{m}}(T_2, v), \nabla T_2' \rangle_{\tilde{m}} &= a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho})(a\partial_a v, \partial_\xi v) + a\Theta_1(\tilde{h})(\tilde{\partial}v, v)) \\ &\quad + a^{n-4} b^{n-1} (\Theta_2(\tilde{h}_{\rho\rho})(a\partial_a v, \partial_\xi v) + a\Theta_2(\tilde{h})(\tilde{\partial}v, \tilde{\partial}v)), \\ \langle \mathcal{F}_{\tilde{g}}(T_2, v), \nabla T_2'' \rangle_{\tilde{g}} - \langle \mathcal{F}_{\tilde{m}}(T_2, v), \nabla T_2'' \rangle_{\tilde{m}} &= a^{\frac{n-7}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho})(a\partial_a v, \partial_\xi v) + a\Theta_1(\tilde{h})(\tilde{\partial}v, v)) \\ &\quad + a^{n-5} b^{n-1} (\Theta_2(\tilde{h}_{\rho\rho})(a\partial_a v, \partial_\xi v) + a\Theta_2(\tilde{h})(\tilde{\partial}v, \tilde{\partial}v)), \\ \mathcal{Q}_{\tilde{g}}(T_2, v) - \mathcal{Q}_{\tilde{m}}(T_2, v) &= a^{\frac{n-7}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho})(a\partial_a v, \partial_\xi v) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho})(a\partial_a v, \partial_\xi v) \\ &\quad + a\Theta_1(\tilde{h})(\tilde{\partial}v, v) + a\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})(\tilde{\partial}v, v)). \end{aligned}$$

Proof. We can do the similar computation as in the proof of Lemma 5.3 by writing out the metric components in local coordinates (a, ξ, θ) with $\xi = -\log b$ and θ is normal w.r.t. the standard spherical metric $d^2\theta$ at $p \in \tilde{\Omega}_2 \cap \text{Int}(X^2)$:

$$\begin{aligned} \tilde{g} &= \tilde{m} + a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} \begin{bmatrix} \tilde{h}_{\rho\rho} & -a(\tilde{h}_{\rho\rho} - a\tilde{h}_{0\rho}) & a[\Theta_1(\tilde{h})]_{1 \times (n-1)} \\ -a(\tilde{h}_{\rho\rho} - a\tilde{h}_{0\rho}) & a^2(\tilde{h}_{\rho\rho} - 2a\tilde{h}_{0\rho} + a^2\tilde{h}_{00}) & a^2[\Theta_1(\tilde{h})]_{1 \times (n-1)} \\ a[\Theta_1(\tilde{h})]_{(n-1) \times 1} & a^2[\Theta_1(\tilde{h})]_{(n-1) \times 1} & a^2[\Theta_1(\tilde{h})]_{(n-1) \times (n-1)} \end{bmatrix}, \\ \tilde{g}^{-1} &= \tilde{m}^{-1} + a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} \begin{bmatrix} -a^2(\tilde{h}_{\rho\rho} + a\Theta_1(\tilde{h})) & -a(\tilde{h}_{\rho\rho} + a\Theta_1(\tilde{h})) & a^2[\Theta_1(\tilde{h})]_{1 \times (n-1)} \\ -a(\tilde{h}_{\rho\rho} + a\Theta_1(\tilde{h})) & -(\tilde{h}_{\rho\rho} + a^{\frac{n-1}{2}}\Theta_2(\tilde{h})) & a[\Theta_1(\tilde{h})]_{1 \times (n-1)} \\ a^2[\Theta_1(\tilde{h})]_{(n-1) \times 1} & a[\Theta_1(\tilde{h})]_{(n-1) \times 1} & a^2[\Theta_1(\tilde{h})]_{(n-1) \times (n-1)} \end{bmatrix}. \end{aligned}$$

Hence

$$\begin{aligned}
\langle \nabla T_2, \nabla T_2 \rangle_{\tilde{g}} - \langle \nabla T_2, \nabla T_2 \rangle_{\tilde{m}} &= -a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} (\tilde{h}_{\rho\rho} (1 + a^{\delta' + \frac{1}{2}})^2 + a\Theta_1(\tilde{h})), \\
\langle \nabla T'_2, \nabla T'_2 \rangle_{\tilde{g}} - \langle \nabla T'_2, \nabla T'_2 \rangle_{\tilde{m}} &= -a^{\frac{n-1}{2}} b^{\frac{n-1}{2}} (\tilde{h}_{\rho\rho} + a\Theta_1(\tilde{h})), \\
\langle \nabla T_2, \nabla T'_2 \rangle_{\tilde{g}} - \langle \nabla T_2, \nabla T'_2 \rangle_{\tilde{m}} &= -a^{\frac{n-3}{2}} b^{\frac{n-1}{2}} (\tilde{h}_{\rho\rho} (1 + a^{\delta' + \frac{1}{2}}) + a\Theta_1(\tilde{h})), \\
\langle \nabla T_2, \nabla T''_2 \rangle_{\tilde{g}} - \langle \nabla T_2, \nabla T''_2 \rangle_{\tilde{m}} &= -a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} (\tilde{h}_{\rho\rho} (1 + a^{\delta' + \frac{1}{2}}) + a\Theta_1(\tilde{h})), \\
\langle \nabla T_2, \nabla v \rangle_{\tilde{g}} - \langle \nabla T_2, \nabla v \rangle_{\tilde{m}} &= a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} (\tilde{h}_{\rho\rho} (1 + a^{\delta' + \frac{1}{2}}) (a\partial_a v + \partial_\xi v) + a\Theta_1(\tilde{h})(\tilde{\partial} v)), \\
\langle \nabla T'_2, \nabla v \rangle_{\tilde{g}} - \langle \nabla T'_2, \nabla v \rangle_{\tilde{m}} &= a^{\frac{n-3}{2}} b^{\frac{n-1}{2}} (\tilde{h}_{\rho\rho} (a\partial_a v + \partial_\xi v) + a\Theta_1(\tilde{h})(\tilde{\partial} v)), \\
\langle \nabla T''_2, \nabla v \rangle_{\tilde{g}} - \langle \nabla T''_2, \nabla v \rangle_{\tilde{m}} &= a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} (\tilde{h}_{\rho\rho} (a\partial_a v + \partial_\xi v) + a\Theta_1(\tilde{h})(\tilde{\partial} v)), \\
\langle \nabla v, \nabla v \rangle_{\tilde{g}} - \langle \nabla v, \nabla v \rangle_{\tilde{m}} &= -a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} (\tilde{h}_{\rho\rho} (a\partial_a v + \partial_\xi v)^2 + a\Theta_1(\tilde{h})(\tilde{\partial} v, \tilde{\partial} v)).
\end{aligned}$$

Then the estimates of quadratic forms follows directly. For $\mathcal{Q}_g(T_2, v)$, first by the formula of $\square_{\tilde{g}}$ in the proof of Proposition 4.1, we have

$$\square_{\tilde{g}} T_2 - \square_{\tilde{m}} T_2 = -a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} \tilde{h}_{\rho\rho} (1 + (\frac{1}{2} - \delta') a^{\delta' + \frac{1}{2}}) + a^{\frac{n-3}{2}} b^{\frac{n-1}{2}} \Theta_1(\tilde{h}),$$

Hence

$$\begin{aligned}
&(\gamma_0 v^2 - \langle \nabla v, \nabla v \rangle_{\tilde{g}}) \square_{\tilde{g}} T_2 - (\gamma_0 v^2 - \langle \nabla v, \nabla v \rangle_{\tilde{m}}) \square_{\tilde{m}} T_2 \\
&= a^{\frac{n-7}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) (a\partial_a v, \partial_\xi v) + a\Theta_1(\tilde{h})(\tilde{\partial} v, v)).
\end{aligned}$$

And moreover,

$$\begin{aligned}
\Gamma_{aa}^a(\tilde{g}) - \Gamma_{aa}^a(\tilde{m}) &= a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_1(\tilde{h}) + a\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) \\
&\quad + a^{n-4} b^{n-1} (\Theta_2(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_2(\tilde{h}) + a\Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{a\xi}^a(\tilde{g}) - \Gamma_{a\xi}^a(\tilde{m}) &= a^{\frac{n-3}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_1(\tilde{h}) + a\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) \\
&\quad + a^{n-3} b^{n-1} (\Theta_2(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_2(\tilde{h}) + a\Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\xi\xi}^a(\tilde{g}) - \Gamma_{\xi\xi}^a(\tilde{m}) &= a^{\frac{n-1}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_1(\tilde{h}) + a\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) \\
&\quad + a^{n-2} b^{n-1} (\Theta_2(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_2(\tilde{h}) + a\Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{a\theta}^a(\tilde{g}) - \Gamma_{a\theta}^a(\tilde{m}) &= a^{\frac{n-3}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) + a^{n-3} b^{n-1} (\Theta_2(\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\xi\theta}^a(\tilde{g}) - \Gamma_{\xi\theta}^a(\tilde{m}) &= a^{\frac{n-1}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) + a^{n-2} b^{n-1} (\Theta_2(\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\theta\theta}^a(\tilde{g}) - \Gamma_{\theta\theta}^a(\tilde{m}) &= a^{\frac{n-1}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) + a^{n-2} b^{n-1} (\Theta_2(\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{aa}^\xi(\tilde{g}) - \Gamma_{aa}^\xi(\tilde{m}) &= a^{\frac{n-7}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_1(\tilde{h}) + a\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) \\
&\quad + a^{n-5} b^{n-1} (\Theta_2(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_2(\tilde{h}) + a\Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{a\xi}^\xi(\tilde{g}) - \Gamma_{a\xi}^\xi(\tilde{m}) &= a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_1(\tilde{h}) + a\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) \\
&\quad + a^{n-4} b^{n-1} (\Theta_2(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_2(\tilde{h}) + a\Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\xi\xi}^\xi(\tilde{g}) - \Gamma_{\xi\xi}^\xi(\tilde{m}) &= a^{\frac{n-3}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_1(\tilde{h}) + a\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) \\
&\quad + a^{n-3} b^{n-1} (\Theta_2(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho}) + a\Theta_2(\tilde{h}) + a\Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{a\theta}^\xi(\tilde{g}) - \Gamma_{a\theta}^\xi(\tilde{m}) &= a^{\frac{n-5}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) + a^{n-4} b^{n-1} (\Theta_2(\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\xi\theta}^\xi(\tilde{g}) - \Gamma_{\xi\theta}^\xi(\tilde{m}) &= a^{\frac{n-3}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) + a^{n-3} b^{n-1} (\Theta_2(\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\theta\theta}^\xi(\tilde{g}) - \Gamma_{\theta\theta}^\xi(\tilde{m}) &= a^{\frac{n-3}{2}} b^{\frac{n-1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})) + a^{n-3} b^{n-1} (\Theta_2(\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h})).
\end{aligned}$$

which implies that

$$\begin{aligned} & \nabla^2 T_2(dv, dv)_{\tilde{g}} - \nabla^2 T_2(dv, dv)_{\tilde{m}} \\ &= a^{\frac{n-7}{2}} b^{\frac{n-1}{2}} ((\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho}))(a\partial_a v, \partial_\xi v) + a(\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h}))(\tilde{\partial}v, \tilde{\partial}v)) \\ & \quad + a^{n-5} b^{n-1} ((\Theta_2(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho}))(a\partial_a v, \partial_\xi v) + a(\Theta_2(\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h}))(\tilde{\partial}v, \tilde{\partial}v)). \end{aligned}$$

We prove the third formula. $\square$

5.3. **In $\tilde{\Omega}_3$.** In $\Omega_3 \cup \Omega_0$, define for $\tau_0 \geq 8$

$$\begin{aligned} T_3 &= t - \psi_2(r), \quad T'_3 = \frac{t - 2\tau_0}{\psi_3(r)} - 1; \\ \tilde{\Omega}_3 &= \overline{\{T_3 \leq \tau_0\} \setminus (\tilde{\Omega}_1 \cup \tilde{\Omega}_2)}, \quad \Sigma_3 = \{T'_3 = -1\} \cap \tilde{\Omega}_3, \quad \Sigma_4 = \{T_3 = \tau_0\}, \end{aligned}$$

where $\psi_2, \psi_3 \in C^2(\tilde{\Omega}_3)$ are defined by

$$\begin{aligned} \psi_2(r) &= \begin{cases} r + \frac{2}{2\delta'+1} r^{-\delta'-\frac{1}{2}} & \text{if } r > 1 \\ 1 + \frac{2}{2\delta'+1} + (\delta' + \frac{3}{2})(\frac{1}{4} - \frac{1}{4}r^2 + \frac{1}{8}r^4) & \text{if } r \leq 1 \end{cases}, \\ \psi_3(r) &= \begin{cases} r & \text{if } r > 1 \\ \frac{3}{8} + \frac{3}{4}r^2 - \frac{1}{8}r^4 & \text{if } r \leq 1 \end{cases}. \end{aligned}$$

Hence $-1 \leq \psi'_2 \leq 1, 0 \leq \psi'_3 \leq 1$. Then in $\tilde{\Omega}_3$

$$\langle \nabla T_3, \nabla T_3 \rangle_{\tilde{m}} = \tilde{\rho}^{-2}(-1 + (\psi'_2(r))^2) < 0.$$

We divide $\tilde{\Omega}_3$ into two parts:

$$\tilde{\Omega}'_3 = \tilde{\Omega}_3 \cap \{-1 \leq T'_3 \leq 0\}, \quad \tilde{\Omega}''_3 = \tilde{\Omega}_3 \cap \{T'_3 \leq -1\}.$$

where $\tilde{\Omega}''_3 \subset \dot{X}$ is compact and $\tilde{\Omega}'_3 \subset \Omega_3$. Recall that in Ω_3 we use the coordinates

$$(\tau, \rho, \theta) = (t - r, \frac{1}{r}, \frac{x}{r}), \quad \tilde{\rho} = \rho_1 = \rho, \quad \rho_0 = \rho_2 = 1.$$

Hence in $\tilde{\Omega}'_3$,

$$\begin{aligned} T_3 &= \tau - \frac{2}{2\delta'+1} \rho^{\delta'+\frac{1}{2}}, \quad T'_3 = -\rho(2\tau_0 - \tau); \\ \langle \nabla T_3, \nabla T_3 \rangle_{\tilde{m}} &= -\rho^{\delta'-\frac{1}{2}}(2 - \rho^{\delta'+\frac{3}{2}}), \\ \langle \nabla T'_3, \nabla T'_3 \rangle_{\tilde{m}} &= -\rho(2\tau_0 - \tau)(2 - \rho(2\tau_0 - \tau)). \end{aligned}$$

Here T_3 is time-like functions all over $\tilde{\Omega}_3 \cap \dot{X}$; T'_3 is time-like in the interior of $\tilde{\Omega}'_3$ and null on $\tilde{\Omega}_3 \cap S_1$. With $\gamma_0 < 0$ a constant, we have the following two lemmas.

Lemma 5.7. In $\tilde{\Omega}'_3$,

$$\begin{aligned} \langle \mathcal{F}_{\tilde{m}}(T_3, v), \nabla T_3 \rangle_{\tilde{m}} &= \frac{1}{2} \rho^{2\delta'-1} (|\partial_\tau v|^2 + |\partial_\tau v + \rho^2 \partial_\rho v|^2) + (1 - \rho^{\delta'+\frac{3}{2}}) |\partial_\rho v|^2 \\ & \quad + \rho^{\delta'-\frac{1}{2}} (1 - \frac{1}{2} \rho^{\delta'+\frac{3}{2}}) (|\partial_\theta v|^2 - \gamma_0 |v|^2), \\ \langle \mathcal{F}_{\tilde{m}}(T_3, v), \nabla T'_3 \rangle_{\tilde{m}} &= \frac{1}{2} \rho^{\delta'-\frac{1}{2}} (2\tau_0 - \tau) (|\partial_\tau v|^2 + |\partial_\tau v + \rho^2 \partial_\rho v|^2) + (1 - \frac{1}{2} \rho(2\tau_0 - \tau) - \frac{1}{2} \rho^{\delta'+\frac{3}{2}}) \rho |\partial_\rho v|^2 \\ & \quad + \frac{1}{2} (2\tau_0 - \tau + \rho^{\delta'+\frac{1}{2}} (1 - \rho(2\tau_0 - \tau))) (|\partial_\theta v|^2 - \gamma_0 |v|^2), \\ \mathcal{Q}_{\tilde{m}}(T_3, v) &= \frac{1}{2} (\frac{1}{2} - \delta') \rho^{\delta'-\frac{3}{2}} (|\partial_\tau v|^2 + |\partial_\tau v + \rho^2 \partial_\rho v|^2) - \rho |\partial_\rho v|^2 + \frac{1}{2} (\delta' + \frac{3}{2}) \rho^{\delta'+\frac{1}{2}} (|\partial_\theta v|^2 - \gamma_0 |v|^2). \end{aligned}$$

Proof. In $\tilde{\Omega}'_3$ under coordinates (τ, ρ, θ) with θ normal w.r.t. standard spherical metric at $p \in \tilde{\Omega}'_3$, we can write the metric components of $\tilde{m} = \rho^2 m$ as follows:

$$\tilde{m} = \begin{bmatrix} -\rho^2 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & \mathbb{1}_{n-1} \end{bmatrix}, \quad \tilde{m}^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & \rho^2 & 0 \\ 0 & 0 & \mathbb{1}_{n-1} \end{bmatrix}.$$

Then the quadratic forms follow from

$$\begin{aligned}\langle \nabla T_3, \nabla T'_3 \rangle_{\tilde{m}} &= -(2\tau_0 - \tau) - \rho^{\delta' + \frac{1}{2}}(1 - \rho(2\tau_0 - \tau)), \\ \langle \nabla T_3, \nabla v \rangle_{\tilde{m}} &= \partial_\rho v - \rho^{\delta' - \frac{1}{2}}(\partial_\tau v + \rho^2 \partial_\rho v), \\ \langle \nabla T'_3, \nabla v \rangle_{\tilde{m}} &= \rho \partial_\rho v - (2\tau_0 - \tau)(\partial_\tau v + \rho^2 \partial_\rho v), \\ \langle \nabla v, \nabla v \rangle_{\tilde{m}} &= 2\partial_\tau v \partial_\rho v + |\rho \partial_\rho v|^2 + |\partial_\theta v|^2.\end{aligned}$$

For $\mathcal{Q}_{\tilde{m}}(T_3, v)$, we have

$$\square_{\tilde{m}} T_3 = -(\frac{3}{2} + \delta') \rho^{\delta' + \frac{1}{2}},$$

and the non-zero connection components at p are

$$\Gamma_{\tau\tau}^\tau(\tilde{m}) = \rho, \quad \Gamma_{\rho\tau}^\rho(\tilde{m}) = -\rho, \quad \Gamma_{\tau\tau}^\rho(\tilde{m}) = \rho^3,$$

which implies that

$$\nabla^2 T(dv, dv) = (\frac{1}{2} - \delta') \rho^{\delta' - \frac{3}{2}} |\partial_\tau v + \rho^2 \partial_\rho v|^2 - \rho |\partial_\rho v|^2 - \rho^{\delta' + \frac{1}{2}} (2\partial_\tau v \partial_\rho v + \rho^2 |\partial_\rho v|^2).$$

Then $\mathcal{Q}_{\tilde{m}}(T_3, v)$ follows. $\square$

Lemma 5.8. For $n \geq 4$ and in $\tilde{\Omega}'_3$,

$$\begin{aligned}\langle \mathcal{F}_{\tilde{g}}(T_3, v), \nabla T'_3 \rangle_{\tilde{g}} - \langle \mathcal{F}_{\tilde{m}}(T_3, v), \nabla T'_3 \rangle_{\tilde{m}} &= \rho^{\frac{n-5}{2}} (\Theta_1(\tilde{h}_{\rho\rho})(\partial_\tau v, \rho \partial_\rho v) + \rho \Theta_1(\tilde{h})(\tilde{\partial} v, v)), \\ \mathcal{Q}_{\tilde{g}}(T_3, v) - \mathcal{Q}_{\tilde{m}}(T_3, v) &= \rho^{\frac{n-7}{2}} ((\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}))(\partial_\tau v, \rho \partial_\rho v) + \rho(\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h}))(\tilde{\partial} v, v)).\end{aligned}$$

Proof. We can do the similar computation as in the proof of Lemma 5.3 by writing out the metric components in local coordinates (τ, ρ, θ) with θ normal w.r.t. the standard spherical metric at $p \in \tilde{\Omega}'_3$:

$$\begin{aligned}\tilde{g} &= \tilde{m} + \rho^{\frac{n-5}{2}} \begin{bmatrix} \rho^4 \Theta_1(\tilde{h}) & \rho^2 \Theta_1(\tilde{h}) & \rho^3 [\Theta_1(\tilde{h})]_{1 \times (n-1)} \\ \rho^2 \Theta_1(\tilde{h}) & \tilde{h}_{\rho\rho} & \rho [\Theta_1(\tilde{h})]_{1 \times (n-1)} \\ \rho^3 [\Theta_1(\tilde{h})]_{(n-1) \times 1} & \rho [\Theta_1(\tilde{h})]_{(n-1) \times (n-1)} & \rho^2 [\Theta_1(\tilde{h})]_{(n-1) \times (n-1)} \end{bmatrix}, \\ \tilde{g}^{-1} &= \tilde{m}^{-1} + \rho^{\frac{n-5}{2}} \begin{bmatrix} -\tilde{h}_{\rho\rho} + \rho^{\frac{n-1}{2}} \Theta_2(\tilde{h}) & \rho^2 \Theta_1(\tilde{h}) & \rho [\Theta_1(\tilde{h})]_{1 \times (n-1)} \\ \rho^2 \Theta_1(\tilde{h}) & \rho^4 \Theta_1(\tilde{h}) & \rho^3 [\Theta_1(\tilde{h})]_{1 \times (n-1)} \\ \rho [\Theta_1(\tilde{h})]_{(n-1) \times 1} & \rho^3 [\Theta_1(\tilde{h})]_{(n-1) \times (n-1)} & \rho^2 [\Theta_1(\tilde{h})]_{(n-1) \times (n-1)} \end{bmatrix}.\end{aligned}$$

Hence the quadratic form estimates follows by

$$\begin{aligned}\langle \nabla T_3, \nabla T'_3 \rangle_{\tilde{g}} - \langle \nabla T_3, \nabla T'_3 \rangle_{\tilde{m}} &= -\rho^{\frac{n-3}{2}} \tilde{h}_{\rho\rho} + \rho^{\frac{n-1}{2}} \Theta_1(\tilde{h}), \\ \langle \nabla T_3, \nabla v \rangle_{\tilde{g}} - \langle \nabla T_3, \nabla v \rangle_{\tilde{m}} &= -\rho^{\frac{n-5}{2}} \tilde{h}_{\rho\rho} \partial_\tau v + \rho^{\frac{n-3}{2}} \Theta_1(\tilde{h})(\tilde{\partial} v), \\ \langle \nabla T'_3, \nabla v \rangle_{\tilde{g}} - \langle \nabla T'_3, \nabla v \rangle_{\tilde{m}} &= -\rho^{\frac{n-3}{2}} \tilde{h}_{\rho\rho} \partial_\tau v + \rho^{\frac{n-1}{2}} \Theta_1(\tilde{h})(\tilde{\partial} v), \\ \langle \nabla v, \nabla v \rangle_{\tilde{g}} - \langle \nabla T_3, \nabla v \rangle_{\tilde{m}} &= -\rho^{\frac{n-5}{2}} \tilde{h}_{\rho\rho} |\partial_\tau v|^2 + \rho^{\frac{n-3}{2}} \Theta_1(\tilde{h})(\tilde{\partial} v, \tilde{\partial} v),\end{aligned}$$

For $\mathcal{Q}_{\tilde{g}}(T_3, v) - \mathcal{Q}_{\tilde{m}}(T_3, v)$, we first have

$$\square_{\tilde{g}} T_3 - \square_{\tilde{m}} T_3 = \rho^{\frac{n-3}{2}} \Theta_1(\tilde{h}).$$

Hence

$$\begin{aligned}(\gamma_0 v^2 - \langle \nabla v, \nabla v \rangle_{\tilde{g}}) \square_{\tilde{g}} T_3 - (\gamma_0 v^2 - \langle \nabla v, \nabla v \rangle_{\tilde{m}}) \square_{\tilde{m}} T_3 \\ = \rho^{\frac{n-3}{2}} \Theta_1(\tilde{h})(\partial_\tau v)(\partial_\rho v) + \rho^{\frac{n-4}{2} + \delta} \Theta_1(\tilde{h})(\tilde{\partial} v, v) + \rho^{n-4} \Theta_2(\tilde{h})(\tilde{\partial} v, v).\end{aligned}$$

Moreover,

$$\begin{aligned}
\Gamma_{\rho\rho}^\rho(\tilde{g}) - \Gamma_{\rho\rho}^\rho(\tilde{m}) &= \rho^{\frac{n-5}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + \rho\Theta_1(\tilde{h}) + \rho\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\rho\tau}^\rho(\tilde{g}) - \Gamma_{\rho\tau}^\tau(\tilde{m}) &= \rho^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + \rho\Theta_1(\tilde{h}) + \rho\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\tau\tau}^\rho(\tilde{g}) - \Gamma_{\tau\tau}^\tau(\tilde{m}) &= \rho^{\frac{n+3}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\rho\theta}^\rho(\tilde{g}) - \Gamma_{\rho\theta}^\tau(\tilde{m}) &= \rho^{\frac{n-3}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\tau\theta}^\rho(\tilde{g}) - \Gamma_{\tau\theta}^\tau(\tilde{m}) &= \rho^{\frac{n+1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\theta\theta}^\rho(\tilde{g}) - \Gamma_{\theta\theta}^\tau(\tilde{m}) &= \rho^{\frac{n-1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\rho\rho}^\tau(\tilde{g}) - \Gamma_{\rho\rho}^\tau(\tilde{m}) &= \rho^{\frac{n-7}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + \rho\Theta_1(\tilde{h}) + \rho\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\rho\tau}^\tau(\tilde{g}) - \Gamma_{\rho\tau}^\tau(\tilde{m}) &= \rho^{\frac{n-5}{2}} (\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}) + \rho\Theta_1(\tilde{h}) + \rho\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\tau\tau}^\tau(\tilde{g}) - \Gamma_{\tau\tau}^\tau(\tilde{m}) &= \rho^{\frac{n-1}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\rho\theta}^\tau(\tilde{g}) - \Gamma_{\rho\theta}^\tau(\tilde{m}) &= \rho^{\frac{n-5}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\tau\theta}^\tau(\tilde{g}) - \Gamma_{\tau\theta}^\tau(\tilde{m}) &= \rho^{\frac{n-3}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})), \\
\Gamma_{\theta\theta}^\tau(\tilde{g}) - \Gamma_{\theta\theta}^\tau(\tilde{m}) &= \rho^{\frac{n-3}{2}} (\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})),
\end{aligned}$$

Hence

$$\begin{aligned}
&\nabla^2 T_3(dv, dv)_{\tilde{g}} - \nabla^2 T_3(dv, dv)_{\tilde{m}} \\
&= \rho^{\frac{n-7}{2}} ((\Theta_1(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{\partial}\tilde{h}_{\rho\rho}))(\partial_\tau v, \rho\partial_\rho v) + \rho(\Theta_1(\tilde{h}) + \Theta_0(\tilde{h})(\tilde{\partial}\tilde{h}))(\tilde{\partial}v, v)) \\
&\quad + \rho^{n-5} ((\Theta_2(\tilde{h}_{\rho\rho}) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho}))(\partial_\tau v, \rho\partial_\rho v) + \rho(\Theta_2(\tilde{h}) + \Theta_1(\tilde{h})(\tilde{\partial}\tilde{h}))(\tilde{\partial}v, v)).
\end{aligned}$$

Then $\mathcal{Q}_{\tilde{g}}(T_3, v) - \mathcal{Q}_{\tilde{m}}(T_3, v)$ follows. $\square$

5.4. **In $\tilde{\Omega}_4$.** In Ω_4 , recall that we have the coordinates and boundary defining functions

$$(\bar{a}, \bar{b}, \theta) = (1 - \frac{r}{t}, \frac{1}{t - r}, \frac{x}{r}); \quad \rho_0 = 1, \quad \rho_1 = \bar{a}, \quad \rho_2 = \bar{b}, \quad \tilde{\rho} = \bar{a}\bar{b};$$

and in Ω_5 , we have

$$(\phi, y) = (\frac{1}{t}, \frac{x}{t}), \quad R = |y|; \quad \rho_0 = \rho_1 = 1, \quad \rho_2 = \tilde{\rho} = \phi.$$

Both sets of coordinates can be extended to $\Omega_4 \cup \Omega_5$ with transition map

$$\phi = \bar{a}\bar{b}, \quad y = (1 - \bar{a})\theta.$$

Define

$$\begin{aligned}
T_4 &= -\frac{2}{2\delta'+1} \left(\frac{\bar{a}(2-\bar{a})}{2}\right)^{\delta'+\frac{1}{2}} + \ln(2-\bar{a}) - \ln \bar{b} = -\frac{2}{2\delta'+1} \left(\frac{1-|y|^2}{2}\right)^{\delta'+\frac{1}{2}} + \ln(1-|y|^2) - \ln \phi, \\
T'_4 &= -\left(\frac{2-\bar{a}}{2}\right)^{1-\alpha} \bar{a}\bar{b}^\alpha = -\left(\frac{1-|y|^2}{2}\right)^{1-\alpha} \phi^\alpha;
\end{aligned}$$

and let

$$\tilde{\Omega}_4 = \overline{(\Omega_4 \cup \Omega_5)} \setminus \tilde{\Omega}_3.$$

Then in $\tilde{\Omega}_4 \cap \mathring{X}$,

$$\begin{aligned}
\langle \nabla T_4, \nabla T_4 \rangle_{\tilde{m}} &= -1 - 2(1-\bar{a})^2 \left(\frac{\bar{a}(2-\bar{a})}{2}\right)^{\delta'-\frac{1}{2}} [1 - \left(\frac{\bar{a}(2-\bar{a})}{2}\right)^{\delta'+\frac{1}{2}}] < 0, \\
\langle \nabla T'_4, \nabla T'_4 \rangle_{\tilde{m}} &= -2\bar{a} \left(\frac{2-\bar{a}}{2}\right)^{2-2\alpha} \bar{b}^{2\alpha} \left(\frac{2-\bar{a}}{2}\alpha - 1 + \bar{a}\right) \left(1 + \frac{\alpha-1}{2-\bar{a}}\bar{a}\right) < 0, \\
\langle \nabla T'_4, \nabla T_4 \rangle_{\tilde{m}} &= -\left(\frac{2-\bar{a}}{2}\right)^{1-\alpha} \bar{b}^\alpha \left(\alpha\bar{a} + \frac{2(1-\bar{a})^2}{2-\bar{a}} (1 - (2-\alpha)\left(\frac{\bar{a}(2-\bar{a})}{2}\right)^{\delta'+\frac{1}{2}})\right) < 0.
\end{aligned}$$

Hence T_4, T'_4 are time-like in $\tilde{\Omega}_4 \cap \mathring{X}$ and T'_4 is null on $S_1^+ \cap \tilde{\Omega}_4$. With $\gamma_0 < 0$ a constant, we have the following two lemmas.

Lemma 5.9. In $\tilde{\Omega}_4$,

$$\begin{aligned} \langle \mathcal{F}_{\tilde{m}}(T_4, v), \nabla T_4' \rangle_{\tilde{m}} &= \left(\frac{2-\bar{a}}{2}\right)^{1-\alpha} \bar{b}^\alpha \left(C_1 (|\bar{a}(2-\bar{a})\partial_{\bar{a}}v - \bar{b}\partial_{\bar{b}}v|^2 + |\bar{b}\partial_{\bar{b}}v|^2) + C_2 |\partial_{\bar{a}}v|^2 \right. \\ &\quad \left. + C_3 \left(\frac{|\partial_{\theta}v|^2}{(1-\bar{a})^2} - \gamma_0 v^2 \right) \right), \\ \bar{a}\mathcal{Q}_{\tilde{m}}(T_4, v) &= D_1 (|\bar{a}(2-\bar{a})\partial_{\bar{a}}v - \bar{b}\partial_{\bar{b}}v|^2 + |\bar{b}\partial_{\bar{b}}v|^2) + D_2 |\partial_{\bar{a}}v|^2 \\ &\quad + D_3 (|\bar{a}(2-\bar{a})\partial_{\bar{a}}v - \bar{b}\partial_{\bar{b}}v|^2 - |\bar{b}\partial_{\bar{b}}v|^2) + D_4 \frac{|\partial_{\theta}v|^2}{(1-\bar{a})^2} + D_5 (-\gamma_0 v^2), \end{aligned}$$

where

$$\begin{aligned} C_1 &= \frac{(1-\bar{a})^2}{2-\bar{a}} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'-\frac{1}{2}} + \frac{1-\bar{a}}{(2-\bar{a})^2} + \frac{\alpha}{(2-\bar{a})^2} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}} + \frac{\alpha\bar{a}}{(2-\bar{a})^2} \left(\frac{1}{2} - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}} \right), \\ C_2 &= \alpha\bar{a}(1-\bar{a}) \left(1 - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}} \right) - \bar{a}(1-\bar{a}), \\ C_3 &= \frac{\alpha\bar{a}}{2} + \frac{(1-\bar{a})^2}{2-\bar{a}} + \frac{(\alpha-2)(1-\bar{a})^2}{2-\bar{a}} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}}; \\ D_1 &= \left(\frac{1}{2} - \delta' \right) \frac{(1-\bar{a})^2}{2-\bar{a}} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'-\frac{1}{2}} - \frac{\bar{a}}{2(2-\bar{a})^2} + \frac{1}{2-\bar{a}} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}}, \\ D_2 &= -\bar{a}(1-\bar{a}), \\ D_3 &= \frac{n-1}{2-\bar{a}} \left(\frac{1}{2} - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}} \right), \\ D_4 &= (1+2\delta') \frac{(1-\bar{a})^2}{2-\bar{a}} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}} + (n-2)\bar{a} \left(\frac{1}{2} - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}} \right), \\ D_5 &= (1+2\delta') \frac{(1-\bar{a})^2}{2-\bar{a}} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}} + n\bar{a} \left(\frac{1}{2} - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}} \right). \end{aligned}$$

Proof. Under coordinates $(\bar{a}, \xi, \theta)$ with $\xi = -\log \bar{b}$ and θ normal w.r.t. the standard spherical metric $d^2\theta$ at $p \in \tilde{\Omega}_4 \cap \text{Int}(X^2)$, the metric components are:

$$\tilde{m} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & -\bar{a}(2-\bar{a}) & 0 \\ 0 & 0 & (1-\bar{a})^2 \mathbb{1}_{n-1} \end{bmatrix}, \quad \tilde{m}^{-1} = \begin{bmatrix} \bar{a}(2-\bar{a}) & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & \frac{1}{(1-\bar{a})^2} \mathbb{1}_{n-1} \end{bmatrix}.$$

Then the quadratic forms follow from

$$\begin{aligned} \langle \nabla T_4, \nabla T_4' \rangle_{\tilde{m}} &= -\bar{b}^\alpha \left(\frac{2-\bar{a}}{2} \right)^{-\alpha} \left(1 - (2-\alpha) \left[\frac{\bar{a}(2-\bar{a})}{2} + (1-\bar{a})^2 \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}} \right] \right), \\ \langle \nabla T_4, \nabla T_4'' \rangle_{\tilde{m}} &= -1 - (1-\bar{a})^2 \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'-\frac{1}{2}}, \\ \langle \nabla T_4, \nabla v \rangle_{\tilde{m}} &= - \left(\left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'-\frac{1}{2}} + \frac{1}{2-\bar{a}} \right) \partial_{\xi} v + (1-\bar{a} - 2 \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}}) \partial_{\bar{a}} v, \\ \langle \nabla T_4', \nabla v \rangle_{\tilde{m}} &= \bar{b}^\alpha \left(\frac{2-\bar{a}}{2} \right)^{-\alpha} \left(- \left(\frac{2-\bar{a}}{2} + \frac{\alpha-1}{2} \bar{a} \right) \partial_{\xi} v + \frac{\alpha-2}{2} (1-\bar{a})(2-\bar{a}) \bar{a} \partial_{\bar{a}} v \right), \\ \langle \nabla T_4'', \nabla v \rangle_{\tilde{m}} &= - \frac{1}{2-\bar{a}} \partial_{\xi} v + (1-\bar{a}) \partial_{\bar{a}} v, \\ \langle \nabla v, \nabla v \rangle_{\tilde{m}} &= 2 \partial_{\bar{a}} v \partial_{\xi} v + \bar{a}(2-\bar{a}) \partial_{\bar{a}} v + \frac{|\partial_{\theta}v|^2}{(1-\bar{a})^2}. \end{aligned}$$

For $\mathcal{Q}_{\tilde{m}}(T_4, v)$, we have

$$\square_{\tilde{m}} T_4 = -(1+2\delta')(1-\bar{a})^2 \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'-\frac{1}{2}} - n + 2n \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta'+\frac{1}{2}}$$

and the non-zero connection components at p are

$$\begin{aligned} \Gamma_{\bar{a}\xi}^{\bar{a}} &= -(1-\bar{a}), \quad \Gamma_{\xi\xi}^{\bar{a}} = \bar{a}(1-\bar{a})(2-\bar{a}), \quad \Gamma_{\theta\theta}^{\bar{a}} = \bar{a}(1-\bar{a})(2-\bar{a}), \\ \Gamma_{\xi\xi}^{\xi} &= 1-\bar{a}, \quad \Gamma_{\theta\theta}^{\xi} = 1-\bar{a}, \end{aligned}$$

which implies $\mathcal{Q}_{\tilde{m}}(T_4, v)$. □

Lemma 5.10. For $n \geq 4$ and $\lambda \in (\frac{1}{2} - \delta', 1 - 2\delta')$, there exists $\alpha \in (1, \frac{n-1}{\lambda})$ such that in $\tilde{\Omega}_4$,

$$|T_4'| \mathcal{Q}_{\tilde{m}}(T_4, v) \leq \lambda \langle \mathcal{F}_{\tilde{m}}(T_4, v), \nabla T_4' \rangle_{\tilde{m}}.$$

Proof. We only need to find $\alpha' = \lambda\alpha \in (1, n-1)$ such that

$$(5.2) \quad \begin{aligned} D_4 \frac{|\partial_{\theta}v|^2}{(1-\bar{a})^2} + D_5 (-\gamma_0 v^2) &\leq \lambda C_3 \left(\frac{|\partial_{\theta}v|^2}{(1-\bar{a})^2} - \gamma_0 v^2 \right), \\ D_3 (|\bar{a}(2-\bar{a})\partial_{\bar{a}}v - \bar{b}\partial_{\bar{b}}v|^2 - |\bar{b}\partial_{\bar{b}}v|^2) &\leq E_1 (\bar{a}(2-\bar{a})\partial_{\bar{a}}v - \bar{b}\partial_{\bar{b}}v|^2 + |\bar{b}\partial_{\bar{b}}v|^2) + E_2 |\partial_{\bar{a}}v|^2, \end{aligned}$$

where

$$\begin{aligned} E_1 &= \lambda C_1 - D_1 = (\lambda - \frac{1}{2} + \delta') \frac{(1-\bar{a})^2}{2-\bar{a}} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' - \frac{1}{2}} + \frac{\bar{a} + 2\lambda(1-\bar{a})}{2(2-\bar{a})^2} \\ &\quad + \frac{\alpha' - 2 + \bar{a}}{(2-\bar{a})^2} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} + \frac{\alpha' \bar{a}}{(2-\bar{a})^2} \left(\frac{1}{2} - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \right) \\ E_2 &= \lambda C_2 - D_2 = \alpha' \bar{a} (1 - \bar{a}) \left(1 - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \right) + (1 - \lambda) \bar{a} (1 - \bar{a}). \end{aligned}$$

First, if $\alpha' \in (n-2, n-1)$, then

$$\begin{aligned} \lambda C_3 - D_4 &= \frac{(\alpha' - n + 2)\bar{a}}{2} + (n-2)\bar{a} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} + \frac{(\alpha' - 2)(1-\bar{a})^2}{2-\bar{a}} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \\ &\quad + \frac{(1-\bar{a})^2}{2-\bar{a}} [\lambda + (1 - 2\delta' - 2\lambda) \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}}] \\ &\geq \lambda \frac{(1-\bar{a})^2}{2-\bar{a}} [1 - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}}] \geq 0. \end{aligned}$$

Similarly, if $\alpha' \in (n-1 - \frac{\lambda}{8}, n-1)$,

$$\begin{aligned} \lambda C_3 - D_5 &= \frac{(\alpha' - n)\bar{a}}{2} + n\bar{a} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} + \frac{(1-\bar{a})^2}{2-\bar{a}} (\alpha' - 2) \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \\ &\quad + \frac{(1-\bar{a})^2}{2-\bar{a}} [\lambda + (1 - 2\delta' - 2\lambda) \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}}] \\ &\geq \frac{\bar{a}}{2} [\alpha' - n + n\bar{a}(2 - \bar{a}) + (\alpha' - 2)(1 - \bar{a})^2] + \lambda \frac{(1-\bar{a})^2}{2-\bar{a}} [1 - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}}] \\ &\geq \frac{\bar{a}(1-\bar{a})^2}{2} [2\alpha' - n - 2 + \frac{\lambda}{4}] \geq 0. \end{aligned}$$

Hence we prove the first inequality in (5.2) for $\alpha' \in (n-1 - \frac{\lambda}{8}, n-1)$. For the second one, there are two cases. First, if $\left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \geq \frac{1}{2}$, then $D_3 \leq 0$. So we only need to show $-D_3 \leq E_1$. In this case, notice that $0 \leq 1 - \bar{a} \leq \frac{\sqrt{2}}{2}$. Let $\alpha' \in (n - \frac{3}{2}, n-1)$. Then

$$\begin{aligned} E_1 + D_3 &\geq \frac{n + \alpha' + (n-2-\alpha'+2\lambda)(1-\bar{a})}{2(2-\bar{a})^2} - \frac{n + (n-\alpha')(1-\bar{a})}{(2-\bar{a})^2} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \\ &\geq \frac{n + \alpha' + (n-1-\alpha'-2\delta')(1-\bar{a})}{2(2-\bar{a})^2} - \frac{n + (n-\alpha')(1-\bar{a})}{(2-\bar{a})^2} \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} := f(\delta'). \end{aligned}$$

Here $f'(\delta') \geq 0$ and hence

$$\begin{aligned} E_1 + D_3 &\geq \frac{1}{(2-\bar{a})^2} \left[\frac{n + \alpha' + (n-1-\alpha')(1-\bar{a})}{2} - (n + (n-\alpha')(1-\bar{a})) \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\frac{1}{2}} \right] \\ &\geq \frac{1}{(2-\bar{a})^2} \left[\frac{n + \alpha'}{2} - \frac{n}{\sqrt{2}} - (n - \alpha' - \frac{n-1-\alpha'}{\sqrt{2}})(1-\bar{a}) \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\frac{1}{2}} \right] \\ &\geq \frac{1}{4(2-\bar{a})^2} [(4 - 2\sqrt{2})n - \frac{3\sqrt{2}}{2} - \frac{5}{2}] \geq 0 \end{aligned}$$

for all $n \geq 4$. Second, if $\left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} < \frac{1}{2}$, Then the second inequality is equivalent to

$$(5.3) \quad 2(D_3 - E_1)\bar{a}(2 - \bar{a})\partial_{\bar{a}}v\bar{b}\partial_{\bar{b}}v \leq 2E_1|\bar{b}\partial_{\bar{b}}v|^2 + (E_2 + \bar{a}^2(2 - \bar{a})^2(E_1 - D_3))|\partial_{\bar{a}}v|^2$$

In this case, $E_1 \geq 0$ and

$$\begin{aligned} &E_2 + \bar{a}^2(2 - \bar{a})^2(E_1 - D_3) \\ &\geq \alpha' \bar{a} (1 - \bar{a}) \left(1 - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \right) + (1 - \lambda) \bar{a} (1 - \bar{a}) \\ &\quad + \bar{a}^2 \left[(\alpha' - 2 + \bar{a}) \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} + \frac{\bar{a} + 2\lambda(1-\bar{a})}{2} + (\alpha' \bar{a} - (n-1)(2 - \bar{a})) \left(\frac{1}{2} - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \right) \right] \\ &\geq \bar{a}^2 \left[\alpha' (1 - \bar{a}) \left(1 - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \right) + \frac{2-\bar{a}}{2} + (\alpha' \bar{a} - (n-1)(2 - \bar{a})) \left(\frac{1}{2} - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \right) \right] \\ &\geq \bar{a}^2 (\alpha' + 2 - n) (2 - \bar{a}) \left(\frac{1}{2} - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \right) \geq 0 \end{aligned}$$

for $\alpha' \in (n-2, n-1)$. Hence (5.3) follows by

$$\begin{aligned} 2E_1(E_2 + \bar{a}^2(2 - \bar{a})^2(E_1 - D_3)) &\geq (D_3 - E_1)^2 \bar{a}^2(2 - \bar{a})^2, \\ \iff E_1(2E_2 + \bar{a}^2(2 - \bar{a})^2 E_1) &\geq \bar{a}^2(2 - \bar{a})^2 D_3^2. \end{aligned}$$

By direct computation,

$$2E_2 + \bar{a}^2(2 - \bar{a})^2 E_1 \geq \alpha' \bar{a} (2 - \bar{a})^2 \left(\frac{1}{2} - \left(\frac{\bar{a}(2-\bar{a})}{2} \right)^{\delta' + \frac{1}{2}} \right)$$

Hence for $\alpha' \in (n - \frac{3}{2}, n - 1)$,

$$\begin{aligned} E_1(2E_2 + \bar{a}^2(2 - \bar{a})^2 E_1) &\geq \alpha'[\alpha' + 1 + (\alpha' - 2 + \bar{a})(2 - \bar{a})]\bar{a}^2(\frac{1}{2} - (\frac{\bar{a}(2 - \bar{a})}{2})^{\delta' + \frac{1}{2}})^2 \\ &\geq \alpha'[2\alpha' - 1]\bar{a}^2(\frac{1}{2} - (\frac{\bar{a}(2 - \bar{a})}{2})^{\delta' + \frac{1}{2}})^2 \\ &\geq \bar{a}^2(2 - \bar{a})^2 D_3^2. \end{aligned}$$

We finish the proof for $\alpha \in (\frac{n-1}{\lambda} - \frac{1}{8}, \frac{n-1}{\lambda})$. $\square$

Lemma 5.11. *For $n \geq 4$ and in $\tilde{\Omega}_4$*

$$\begin{aligned} \langle F(T_4, v), \nabla T_4' \rangle_{\tilde{g}} - \langle F(T_4, v), \nabla T_4' \rangle_{\tilde{m}} &= \bar{a}^{\frac{n-5}{2}} \bar{b}^{\frac{n-1}{2} + \alpha} (\Theta_1(\tilde{h}_{\rho\rho})(\bar{a}\partial_{\bar{a}}v, \partial_{\xi}v) + \bar{a}\Theta_1(\tilde{h})(\tilde{\partial}v, v)) \\ &\quad + \bar{a}^{n-4} \bar{b}^{n-1+\alpha} (\Theta_2(\tilde{h}_{\rho\rho})(\bar{a}\partial_{\bar{a}}v, \partial_{\xi}v) + \bar{a}\Theta_2(\tilde{h})(\tilde{\partial}v, \tilde{\partial}v)), \\ \mathcal{Q}_{\tilde{g}}(T_4, v) - \mathcal{Q}_{\tilde{m}}(T_4, v) &= \bar{a}^{\frac{n-7}{2}} \bar{b}^{\frac{n-1}{2}} (\Theta_1(\tilde{h}_{\rho\rho})(\bar{a}\partial_{\bar{a}}v, \partial_{\xi}v) + \Theta_1(\tilde{h}_{\rho\rho})(\tilde{\partial}\tilde{h}_{\rho\rho})(\bar{a}\partial_{\bar{a}}v, \partial_{\xi}v) \\ &\quad + \bar{a}\Theta_1(\tilde{h})(\tilde{\partial}v, v) + \bar{a}\Theta_0(\tilde{h})(\tilde{\partial}\tilde{h})(\tilde{\partial}v, v)). \end{aligned}$$

Proof. The proof is similar as the proof of Lemma 5.6 by substituting (a, b) by $(\bar{a}, \bar{b})$ if restricting to Ω_4 and similar as the proof of Lemma 5.4 if restricting to Ω_5 . $\square$

6. ENERGY ESTIMATES

In this section, we mainly prove the following theorem by energy estimates.

Theorem 3. *For $n \geq 4$ if given Cauchy data*

$$(6.1) \quad (h^0, h^1) \in \tilde{\mathcal{V}}_{\epsilon}^{N, \delta + \frac{n-1}{2}} \quad \text{with } N > n + 6, \delta \in (0, \frac{1}{2}) \text{ and } \epsilon > 0 \text{ small,}$$

where $\tilde{\mathcal{V}}_{\epsilon}^{N, \delta + \frac{n-1}{2}}$ is defined in Section 3, then there exists a global solution h to the reduced Einstein vacuum equations such that $\tilde{h} = \tilde{\rho}^{-\frac{n-1}{2}} h$ is $C^{0, \delta}$ up to S_1 and hence the radiation field in Friedlander's sense (see [Fr]) is well defined: for some $\sigma > 0$ and $C > 0$

$$(6.2) \quad \mathcal{R}_{\mathcal{F}}(h^0, h^1) := \tilde{h}|_{S_1^{\pm}} \in \rho_0^{\delta} \rho_2^{\sigma - \frac{n-1}{2}} H_b^N(S_1^{\pm}), \quad \|\tilde{h}|_{S_1^{\pm}}\|_{\rho_0^{\delta} \rho_2^{\sigma - \frac{n-1}{2}} H_b^N(S_1^{\pm})} \leq C\epsilon.$$

which provides the leading term of h along the bicharacteristic curves.

The theorem follows by Proposition 6.1, 6.2, 6.3, 6.4 and Corollary 2, 3, 4. We do energy estimates for $\tilde{h}$ satisfying the conformal transformation of reduced Einstein equations (4.5), with particular choice of time-like functions and boundary defining functions in each domain $\tilde{\Omega}_i$ for $1 \leq i \leq 4$. Here N, δ are fixed all over this section and ϵ is chosen to be small enough. The image space of $\mathcal{R}_F$ will be refined in Section 7.

6.1. In $\tilde{\Omega}_1$. To solve the equations (4.5) in $\tilde{\Omega}_1$, it is in fact a local existence theorem. We prove the local existence and uniqueness theorem in the conformal setting for $T_1 \in [0, \frac{3}{4}]$ for Cauchy data in weighted b-Sobolev space and small on Σ_0 . Denote by $\Sigma_1^{\mathfrak{t}} = \{T_1 = \mathfrak{t}\}$ the space-like hypersurface with defining function T_1 and $\Omega_1^{\mathfrak{t}}$ the domain bounded by $\Sigma_0, \Sigma_1^{\mathfrak{t}}$ and S_0 . Define the high energy norm on $\Sigma_1^{\mathfrak{t}}$ by

$$M_1^N(\mathfrak{t}; v, \tilde{g}) = \left(\int_{\Sigma_1^{\mathfrak{t}}} \sum_{|I| \leq N} e^{-2\delta \log \rho_0} \langle F(T_1, \tilde{D}^I v), \nabla T_1 \rangle_{\tilde{g}} d\mu_{\tilde{g}}^{T_1} \right)^{\frac{1}{2}} \quad \forall \mathfrak{t} \in [0, \frac{3}{4}].$$

Here $d\mu_{\tilde{g}}^{T_1} \wedge dT_1 = d\text{vol}_{\tilde{g}}$ and $\tilde{D} \in \chi(r)\mathcal{B}_0 + (1 - \chi(r))\mathcal{B}_1$ where $0 \leq \chi \leq 1$ is a smooth cutoff function such that $\chi(r) = 1$ if $r > 1$ and $\chi(r) = 0$ if $r < \frac{1}{2}$. Here

$$\mathcal{B}_0 = \{\partial_{\mu} : \mu = 0, \dots, n\}, \quad \mathcal{B}_1 = \{Z_{\mu\nu} : \mu, \nu = 0, \dots, n\}.$$

We choose $\gamma_0 = -\frac{(n+1)(n-3)}{4}$ for $\mathcal{F}_{\tilde{m}}(T_1, v)$ in Definition 5. First, (6.1) implies that for some constant $C_0 > 0$,

$$(6.3) \quad M_1^N(0; \tilde{h}, \tilde{m}) < C_0 \epsilon.$$

Lemma 6.1. *There exists $\epsilon_1 > 0$ such that if $\|\rho_0^{\frac{n-1}{2}} \tilde{h}\|_{L^\infty(\Sigma_1^t)} < \epsilon_1$ then*

$$\begin{aligned} \frac{1}{C} \|v\|_{\rho_0^\delta H_b^N(\Sigma_1^t)} &\leq \frac{1}{\sqrt{2}} M_1^N(\mathbf{t}; v, \tilde{m}) \leq M_1^N(\mathbf{t}; v, \tilde{g}) \leq \sqrt{2} M_1^N(\mathbf{t}; v, \tilde{m}) \leq C \|v\|_{\rho_0^\delta H_b^N(\Sigma_1^t)}, \\ \sum_{|I| \leq \frac{N}{2}+1} |\tilde{D}^I v| &\leq C \rho_0^\delta M_1^N(\mathbf{t}; v, \tilde{g}), \end{aligned}$$

for some constant C which is independent of ϵ_1 if it is small enough.

We find out the existence of h to (1.3) (resp. $\tilde{h}$ to (4.5)) by considering it as the limit of a sequence h^l (resp. $\tilde{h}^l$), which are solutions to the linear equations (resp. their conformal transformation) for $l \geq 0$ starting with $h^{-1} = 0$ (resp. $\tilde{h}^{-1} = 0$):

$$\begin{aligned} \square_{g^l} h_{\mu\nu}^{l+1} &= F_{\mu\nu}(h^l)(\partial h^l, \partial h^{l+1}), \\ (\square_{\tilde{g}^l} + \gamma(\tilde{h}^l)) \tilde{h}^{l+1} &= \tilde{\rho}_0^{\frac{n-1}{2}} \tilde{F}(\tilde{h}^l, \tilde{h}^{l+1}). \end{aligned}$$

Here $g^l = m + h^l$, $\tilde{g}^l = \tilde{\rho}^2 g^l$ and h^l all have same Cauchy data with (h^0, h^1) . Notice that there is no guarantee that g^l should satisfy the harmonic gauge conditions in coordinates (t, x) . However, it does not change the expression of conformal transformation of equations too much as stated in Section 4.4, i.e.

$$\begin{aligned} \gamma(\tilde{h}^l) &= \gamma_0 + \rho_0^{\frac{n-1}{2}} (\Theta_1(\tilde{h}^l) + \Theta_1(\tilde{\partial} \tilde{h}^l)), \\ \tilde{F}(\tilde{h}^l, \tilde{h}^{l+1}) &= \Theta_0(\tilde{h}^l)(\tilde{\partial} \tilde{h}^l, \tilde{\partial} \tilde{h}^{l+1}) + \Theta_0(\tilde{h}^l)(\tilde{\partial} \tilde{h}^l, \tilde{h}^{l+1}) + \Theta_0(\tilde{h}^l)(\tilde{h}^l, \tilde{\partial} \tilde{h}^{l+1}) + \Theta_0(\tilde{h}^l)(\tilde{h}^l, \tilde{h}^{l+1}). \end{aligned}$$

We also study the equation for $h^{l+1} - h^l$ and $\tilde{h}^{l+1} - \tilde{h}^l$ to get the convergence of h^l and $\tilde{h}^l$:

$$\begin{aligned} (6.4) \quad \square_{g^l} (h^{l+1} - h^l) &= (\square_{g^{l-1}} - \square_{g^l}) h^l + (F(\partial h^l, \partial h^{l+1}) - F(\partial h^{l-1}, \partial h^l)) \\ (\square_{\tilde{g}^l} + \gamma(\tilde{h}^l)) (\tilde{h}^{l+1} - \tilde{h}^l) &= (\square_{\tilde{g}^{l-1}} - \square_{\tilde{g}^l}) \tilde{h}^l + (\gamma(\tilde{h}^{l-1}) - \gamma(\tilde{h}^l)) \tilde{h}^l \\ &\quad + \tilde{\rho}_0^{\frac{n-1}{2}} (\tilde{F}(\tilde{h}^l, \tilde{h}^{l+1}) - \tilde{F}(\tilde{h}^{l-1}, \tilde{h}^l)). \end{aligned}$$

Lemma 6.2. *There exists $\epsilon'_1 > 0$ such that if $M_1^N(\mathbf{t}; \tilde{h}^i, \tilde{g}^{i-1}) < \epsilon'_1$ for all $i = 0, \dots, l$ and $\mathbf{t} \in [0, \frac{3}{4}]$ then*

$$M_1^N(\mathbf{t}; \tilde{h}^{l+1}, \tilde{g}^l) \leq C M_1^N(0; \tilde{h}^{l+1}, \tilde{g}^l) \quad \forall \mathbf{t} \in [0, \frac{3}{4}].$$

where C is a constant independent of $\mathbf{t}$ and $M_1^N(0; \tilde{h}^{l+1}, \tilde{g}^l)$ for ϵ'_1 small enough.

Proof. We can first choose ϵ'_1 such that such that Lemma 6.1 holds for all $\tilde{h}^i$ for $i = 0, \dots, l$. This can be done by induction since $\tilde{h}^{-1} = 0$ and l is finite. So

$$\begin{aligned} \sum_{|I| \leq \frac{N}{2}+1} \|\tilde{D}^I \tilde{h}^i\|_{L^\infty(\Sigma_1^t)} &\leq C M_1^N(\mathbf{t}; \tilde{h}^i, \tilde{g}^{i-1}) \leq C \epsilon'_1, \quad \forall i = 0, \dots, l, \\ \sum_{|I| \leq \frac{N}{2}+1} \|\tilde{D}^I \tilde{h}^{l+1}\|_{L^\infty(\Sigma_1^t)} &\leq C M_1^N(\mathbf{t}; \tilde{h}^{l+1}, \tilde{g}^l), \end{aligned}$$

To get the high energy estimate for $\tilde{h}^{l+1}$ we first deduce the equation for $\tilde{D}^I \tilde{h}^{l+1}$ for $|I| = k \leq N$,

$$\begin{aligned} (\square_{\tilde{g}^l} + \gamma(\tilde{h}^l)) \tilde{D}^I \tilde{h}^{l+1} &= \tilde{D}^I (\rho_0^{\frac{n-1}{2}} \tilde{F}(\tilde{h}^l)(\tilde{h}^l, \tilde{h}^{l+1})) + [\square_{\tilde{g}^l} + \gamma(\tilde{h}^l), \tilde{D}^I] \tilde{h}^{l+1} \\ &= \rho_0^{\frac{n-1}{2}} \sum_{|\alpha| \leq k+1, i \leq k+1, |\alpha|+i \leq k+2, 1 \leq j} \Theta_0(\tilde{h}^l)(\tilde{\partial}^{\alpha_1} \tilde{h}^l, \dots, \tilde{\partial}^{\alpha_j} \tilde{h}^l, \tilde{\partial}^i \tilde{h}^{l+1}) \end{aligned}$$

By Lemma 5.2,

$$(M_1^N(\mathbf{t}; \tilde{h}^{l+1}, \tilde{g}^l)^2 - (M_1^N(0; \tilde{h}^{l+1}, \tilde{g}^l))^2) = \int_{\Omega_1^t} \sum_{|I| \leq N} \operatorname{div}_{\tilde{g}^l} (e^{-2\delta \log \rho_0} F(T_1, \tilde{D}^I \tilde{h})) d\operatorname{vol}_{\tilde{g}^l}.$$

Here by Lemma 5.3 and 5.4

$$\begin{aligned}
& \sum_{|I| \leq N} \operatorname{div}_{\tilde{g}^l} (e^{-2\delta \log \rho_0} \mathcal{F}_{\tilde{g}^l}(T_1, \tilde{D}^I \tilde{h}^{l+1})) \\
&= \sum_{|I| \leq N} e^{-2\delta \log \rho_0} (-2\delta \langle \mathcal{F}_{\tilde{g}^l}(T_1, \tilde{D}^I \tilde{h}^{l+1}), \nabla \log \rho_0 \rangle_{\tilde{g}^l} + \operatorname{div}_{\tilde{g}^l} (\mathcal{F}_{\tilde{g}^l}(T_1, \tilde{D}^I \tilde{h}^{l+1}))) \\
&\leq \sum_{|I| \leq N} e^{-2\delta \log \rho_0} \langle \nabla T_1, \nabla \tilde{D}^I \tilde{h}^{l+1} \rangle_{\tilde{g}^l} (\square_{\tilde{g}^l} + \gamma_0) \tilde{D}^I \tilde{h}^{l+1} + C'(M_1^N(T_1; \tilde{h}^{l+1}, \tilde{g}^l))^2.
\end{aligned}$$

Here ∇ is the covariant derivative w.r.t. $\tilde{g}^l$. Hence

$$\begin{aligned}
\partial_t (M_1^N(\mathbf{t}; \tilde{h}^{l+1}, \tilde{g}^l))^2 &= \int_{\Sigma_1^t} \sum_{|I| \leq N} \operatorname{div}_{\tilde{g}^l} (e^{-2\delta \log \rho_1} \mathcal{F}_{\tilde{g}^l}(T_1, \tilde{D}^I \tilde{h})) d\mu_{\tilde{g}^l}^{T_1} \\
&\leq 2C'(M_1^N(\mathbf{t}; \tilde{h}^l, \tilde{g}^l) + 1)(M_1^N(\mathbf{t}; \tilde{h}^{l+1}, \tilde{g}^l))^2 \\
&\leq C'(1 + 2\epsilon'_1) M_1^N(\mathbf{t}; \tilde{h}^{l+1}, \tilde{g}^l),
\end{aligned}$$

which implies that

$$M_1^N(\mathbf{t}; \tilde{h}^{l+1}, \tilde{g}^l) \leq e^{C'(1+2\epsilon'_1)t} M_1^N(0; \tilde{h}^{l+1}, \tilde{g}^l) < C M_1^N(0; \tilde{h}^{l+1}, \tilde{g}^l), \quad \forall \mathbf{t} \in [0, \frac{3}{4}].$$

Here $C > 0$ is a constant independent of ϵ'_1 if it is small enough. We finish the proof. $\square$

Proposition 6.1. *There exists $\epsilon''_1 > 0$ such that if $M_1^N(0; \tilde{h}, \tilde{m}) < \epsilon''_1$ then the equations (4.5) have a unique solution in $\tilde{\Omega}_1$ such that*

$$M_1^N(\mathbf{t}; \tilde{h}, \tilde{m}) \leq C M_1^N(0; \tilde{h}, \tilde{m}), \quad \forall \mathbf{t} \in [0, \frac{3}{4}],$$

where $C > 0$ is a constant independent of $M_1^N(0; \tilde{h}, \tilde{m})$ if ϵ''_1 is small enough.

Proof. We first choose $\epsilon''_1 > 0$ small such that $M_1^N(\mathbf{t}; \tilde{h}^0, \tilde{g}^{-1}) \leq C\epsilon''_1 < \epsilon'_1$ where $\epsilon'_1 > 0$ small such that Lemma 6.2 holds. This can be done since $\tilde{h}^0$ is the solution to standard wave equation on Minkowski metric and hence can be written out explicitly by using fundamental solution. Then by induction in Lemma 6.2

$$M_1^N(T_1; \tilde{h}^l, \tilde{g}^{l-1}) < C\epsilon''_1 < \epsilon'_1 \quad \forall l \geq 0.$$

We do the similar energy estimates for $\tilde{h}^{l+1} - \tilde{h}^l$ satisfying equations (6.4) and hence

$$\begin{aligned}
(\square_{\tilde{g}^l} + \gamma(\tilde{h}^l)) \tilde{D}^I (\tilde{h}^{l+1} - \tilde{h}^l) &= \tilde{D}^I [\rho_0^{\frac{n-1}{2}} (\tilde{F}(\tilde{h}^l)(\tilde{h}^l, \tilde{h}^{l+1}) - \tilde{F}(\tilde{h}^{l-1})(\tilde{h}^{l-1}, \tilde{h}^l))] \\
&\quad + [\square_{\tilde{g}^l} + \gamma(\tilde{h}^l), \tilde{D}^I] (\tilde{h}^{l+1} - \tilde{h}^l) + D^I [(\square_{\tilde{g}^{l-1}} - \square_{\tilde{g}^l}) \tilde{h}^l + (\gamma(\tilde{h}^{l-1}) - \gamma(\tilde{h}^l)) \tilde{h}^l].
\end{aligned}$$

We deduce the estimates for $\tilde{h}^{l+1} - \tilde{h}^l$ in the same way as in the proof of Lemma 6.2 and have

$$\begin{aligned}
& \partial_t (M_1^N(\mathbf{t}; \tilde{h}^{l+1} - \tilde{h}^l, \tilde{g}^l))^2 \\
&= \int_{\Sigma_1^t} \sum_{|I| \leq N} \operatorname{div}_{\tilde{g}^l} (e^{-2\delta \log \rho_0} \mathcal{F}_{\tilde{g}^l}(T_1, \tilde{D}^I (\tilde{h}^{l+1} - \tilde{h}^l))) d\mu_{\tilde{g}^l}^{T_1} \\
&\leq 2C'(M_1^N(\mathbf{t}; \tilde{h}^{l+1} - \tilde{h}^l, \tilde{g}^l))^2 + 2C'\epsilon'_1 M_1^N(\mathbf{t}; \tilde{h}^l - \tilde{h}^{l-1}, \tilde{g}^l) M_1^N(\mathbf{t}; \tilde{h}^{l+1} - \tilde{h}^l, \tilde{g}^l)
\end{aligned}$$

where $C' > 0$ is a constant independent of ϵ'_1 , and it implies that

$$\begin{aligned}
M_1^N(\mathbf{t}; \tilde{h}^{l+1} - \tilde{h}^l, \tilde{g}^l) &\leq C'\epsilon'_1 e^{C't} \int_0^t e^{-C'T_1} M_1^N(T_1; \tilde{h}^l - \tilde{h}^{l-1}, \tilde{g}^l) dT_1 \\
&\leq 2C'\epsilon'_1 e^{C't} \int_0^t e^{-C'T_1} M_1^N(T_1; \tilde{h}^l - \tilde{h}^{l-1}, \tilde{g}^{l-1}) dT_1.
\end{aligned}$$

For $l \geq 0$, let

$$\mu_l = \sup_{\mathbf{t} \in [0, \frac{3}{4}]} M_1^N(\mathbf{t}; \tilde{h}^l - \tilde{h}^{l-1}, \tilde{m}).$$

Then $\mu_0 \leq \epsilon'_1$ and for $l \geq 1$, $\mu_l \leq 4\epsilon'_1 e^{\frac{3}{4}C} \mu_{l-1}$. Choose ϵ'_1 and hence ϵ''_1 even smaller if necessary such that $4\epsilon'_1 e^{\frac{3}{4}C} \leq \frac{1}{2}$ then for any $\mathbf{t} \in [0, \frac{3}{4}]$, $l, k \geq 1$

$$(M_1^N(\mathbf{t}; \tilde{h}^{l+k}, \tilde{m}))^2 - (M_1^N(\mathbf{t}; \tilde{h}^l, \tilde{m}))^2 = \sum_{i=0}^{k-1} (M_1^N(\mathbf{t}; \tilde{h}^{l+i+1} - \tilde{h}^{l+i}, \tilde{m}))^2 \leq (\frac{1}{4})^l \sum_{i=0}^{\infty} (\frac{1}{4})^i \mu_0^2 \leq (\frac{1}{4})^l \frac{4}{3} \epsilon_1'^2.$$

Hence $\tilde{h}^l$ converges to $\tilde{h}$ with $M_1^N(\mathbf{t}; \tilde{h}, \tilde{m}) \leq 2\epsilon'_1$ for all $\mathbf{t} \in [0, \frac{3}{4}]$. Here $\tilde{h}$ provides a solution to equations (4.5). Choose ϵ'_1 and hence ϵ''_1 small such that Lemma 6.2 holds for $\tilde{h}$. We finish the proof of existence.

For the uniqueness, suppose there is another solutions $\tilde{h}'$ with the same Cauchy data such that $M_1^N(\mathbf{t}; \tilde{h}', \tilde{m}) \leq CM_1^N(\mathbf{t}; \tilde{h}, \tilde{m}) \leq C\epsilon''_1 < \epsilon'_1$ for some constant $C > 0$, then

$$\begin{aligned} (\square_{\tilde{g}} + \gamma(\tilde{h}))(\tilde{h} - \tilde{h}') &= (\square_{\tilde{g}'} - \square_{\tilde{g}})\tilde{h}' + (\gamma(\tilde{h}') - \gamma(\tilde{h}))\tilde{h}' + \rho_0^{\frac{n-1}{2}}(\tilde{F}(\tilde{h})(\tilde{h}, \tilde{h}) - \tilde{F}(\tilde{h}')(\tilde{h}', \tilde{h}')), \\ (\square_{\tilde{g}} + \gamma(\tilde{h}))\tilde{D}^I(\tilde{h} - \tilde{h}') &= \tilde{D}^I(\rho_0^{\frac{n-1}{2}}(\tilde{F}(\tilde{h})(\tilde{h}, \tilde{h}) - \tilde{F}(\tilde{h}')(\tilde{h}', \tilde{h}'))) + [\square_{\tilde{g}'} + \gamma(\tilde{h}'), \tilde{D}^I](\tilde{h}^{l+1} - \tilde{h}^l) \\ &\quad + D^I[(\square_{\tilde{g}'} - \square_{\tilde{g}})\tilde{h}' + (\gamma(\tilde{h}') - \gamma(\tilde{h}))\tilde{h}']. \end{aligned}$$

A similar estimates shows that

$$\begin{aligned} M_1^N(\mathbf{t}; \tilde{h} - \tilde{h}', \tilde{g}) &\leq C\epsilon_1 e^{C\mathbf{t}} \int_0^{\mathbf{t}} e^{-CT_1} M_1^N(T_1; \tilde{h} - \tilde{h}', \tilde{g}) dT_1, \\ \implies \sum_{\mathbf{t} \in [0, \frac{3}{4}]} M_1^N(\mathbf{t}; \tilde{h} - \tilde{h}', \tilde{g}) &\leq \frac{1}{2} \sum_{\mathbf{t} \in [0, \frac{3}{4}]} M_1^N(\mathbf{t}; \tilde{h} - \tilde{h}', \tilde{g}) \end{aligned}$$

Hence $M_1^N(\mathbf{t}; \tilde{h} - \tilde{h}', \tilde{g}) = 0$ for all $\mathbf{t} \in [0, \frac{3}{4}]$, which shows that $\tilde{h} = \tilde{h}'$. $\square$

Hence if $\epsilon > 0$ small enough such that $C_0\epsilon < \epsilon''_1$, we solve the reduced Einstein equations (4.5) up to Σ_1 .

Remark 2. When restricting to $\tilde{\Omega}_1$, all the proofs work for spacial dimension $n \geq 3$.

6.2. In $\tilde{\Omega}_2$. To solve the equations (4.5) up to $\tilde{\Omega}_2$ with given Cauchy data, we first notice that the solution we find in $\tilde{\Omega}_1$ can always be extended a bit over Σ_1 since $\tilde{\Omega}_1$ is closed. If the Cauchy data is small enough, we show that this extension never stop in $\tilde{\Omega}_2$ until it arrives S_1^+ .

For $\mathbf{t} \in [-\frac{1}{4}, 0]$, denote by

$$\tilde{\Sigma}_2^t = \{T_2' = \mathbf{t}\} \cap \tilde{\Omega}_2, \quad \Sigma_2^t = \Sigma_2 \cap \{-\frac{1}{4} \leq T_2' \leq \mathbf{t}\}$$

the space-like hypersurface w.r.t. $\tilde{m}$ with defining function T_2' and T_2 separately, and Ω_2^t the domain bounded by $\Sigma_1, S_0, \Sigma_1^t, \Sigma_2^t$. Define the high energy norm on Σ_1^t and Σ_2^t by

$$\begin{aligned} M_2^N(\mathbf{t}; v, \tilde{g}) &= \left(\int_{\Sigma_2^t} \sum_{|I| \leq N} e^{-2\delta \log \rho_0} \langle \mathcal{F}_{\tilde{g}}(T_2, \tilde{D}^I v), \nabla T_2' \rangle_{\tilde{g}} d\mu_{\tilde{g}}^{T_2'} \right)^{\frac{1}{2}}, \\ L_2^N(\mathbf{t}; v, \tilde{g}) &= \left(\int_{\Sigma_2^t} \sum_{|I| \leq N} e^{-2\delta \log \rho_0} \langle \mathcal{F}_{\tilde{g}}(T_2, \tilde{D}^I v), \nabla T_2 \rangle_{\tilde{g}} d\mu_{\tilde{g}}^{T_2} \right)^{\frac{1}{2}}. \end{aligned}$$

Here $d\mu_{\tilde{g}}^{T_2'} \wedge dT_2' = d\text{vol}_{\tilde{g}}$, $d\mu_{\tilde{g}}^{T_2} \wedge dT_2 = d\text{vol}_{\tilde{g}}$ and $\tilde{D} \in \mathcal{B}_2 = \{Z_{\mu\nu} : \mu, \nu = 0, \dots, n\}$. We choose $\gamma_0 = -\frac{(n-3)(n-1)}{4}$ in the definition of quadratic form $\mathcal{F}_{\tilde{g}}(T_2, v)$. With above notation, first Proposition 6.1 implies that for some $C_1 > 0$

$$M_2^N(-\frac{1}{4}; v, \tilde{g}) < C_1\epsilon, \quad L_2^N(-\frac{1}{4}; v, \tilde{g}) = 0.$$

Recall that we choose $\delta' \in (0, \delta)$ such that $\frac{1}{2} - \delta' < 1 - 2\delta$ in the definition of T_2 .

Lemma 6.3. For $n \geq 4$ there exists $\epsilon_2 > 0$ such that if $\|\rho_0^{\frac{n-1}{2}} \tilde{h}\|_{L^\infty(\Sigma_1^t)} + \|\rho_0^{\frac{n-1}{2}} \rho_1^{-\delta} \tilde{h}_{\rho\rho}\|_{L^\infty(\Sigma_1^t)} < \epsilon_2$, then for $D \in \mathcal{B}_2 = \{Z_{\mu\nu} : \mu\nu = 0, 1, \dots, n\}$,

$$\begin{aligned} \frac{1}{\sqrt{2}} M_2^N(\mathbf{t}; v, \tilde{m}) &\leq M_2^N(\mathbf{t}; v, \tilde{g}) \leq \sqrt{2} M_2^N(\mathbf{t}; v, \tilde{m}), \\ \sum_{|I| \leq \frac{N}{2}+1} |\tilde{D}^I v| &\leq C \rho_0^\delta M_2^N(\mathbf{t}; v, \tilde{g}), \quad \sum_{|I| \leq \frac{N}{2}} |\tilde{D}^I \partial_a v| \leq C \rho_0^\delta \rho_1^{-\frac{1}{2}} M_2^N(\mathbf{t}; v, \tilde{g}), \\ \sum_{|I| \leq \frac{N}{2}} |\tilde{D}^I \tilde{h}_{\rho\rho}| &\leq C \rho_0^\delta \rho_1^{\frac{1}{2}} M_2^N(\mathbf{t}; \tilde{h}, \tilde{m}). \end{aligned}$$

for some constant C independent of ϵ_2 if it is small enough.

Proof. The first three inequality follow from Lemma 5.5 and 5.6 directly and the last one follows from the harmonic gauge conditions. By Lemma 4.3,

$$\begin{aligned} |\tilde{h}_{\rho\rho}| &= \left| \int_0^{\rho_0} b \partial_b \tilde{h}_{\rho\rho} \frac{db}{b} \right| \\ &\leq \int_0^{\rho_0} |a \partial_a \tilde{h}_{\rho\rho} + a \tilde{D} \tilde{h} + (ab)^{\frac{n-1}{2}} \Theta_1(\tilde{h})(\tilde{D} \tilde{h})| \frac{db}{b} \\ &\leq \left(\int_0^{\rho_0} b^\delta \frac{db}{b} \right) a^{\frac{1}{2}} (1 + \Theta_1(\|(ab)^{\frac{n-1}{2}} \tilde{h}\|_\infty)) M_2^N(\mathbf{t}; \tilde{h}, \tilde{m}) \\ &\leq C' \rho_0^\delta \rho_1^{\frac{1}{2}} M_2^N(\mathbf{t}; \tilde{h}, \tilde{m}). \end{aligned}$$

For $|\tilde{D}^I \tilde{h}_{\rho\rho}|$, it is estimated in the same way as $|\tilde{h}_{\rho\rho}|$ since $[b \partial_b, \tilde{D}] = 0$ for all $\tilde{D} \in \mathcal{B}_2$. $\square$

Proposition 6.2. For $n \geq 4$, there exists $\epsilon_2'' > 0$ such that if $M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{m}) \leq \epsilon_2''$, then the equations (4.5) have a unique solution in $\tilde{\Omega}_2$ such that

$$M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) + L_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) \leq C |\mathbf{t}|^{\delta - \frac{1}{2}} M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{g}), \quad \forall \mathbf{t} \in [-\frac{1}{4}, 0],$$

where $C > 0$ is a constant depending on $\delta - \delta'$ but independent of ϵ_2'' if it is small enough.

Proof. By the proof of existence theorem in $\tilde{\Omega}_1$, we see that first we can choose $\epsilon_2'' > 0$ small such that the we can solve the equations (4.5) for $\mathbf{t} < \mathbf{t}_0$, where $\mathbf{t}_0 \in (-\frac{1}{4}, 0)$ such that Lemma 6.3 hold. Then by Lemma 5.2

$$(M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 + (L_2^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 - (M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{g}))^2 = \int_{\Omega_2^t} \sum_{|I| \leq N} \text{div}_{\tilde{g}}(e^{-2\delta \log \rho_0} \mathcal{F}_{\tilde{g}}(T_2, \tilde{D}^I \tilde{h})) d\text{vol}_{\tilde{g}}.$$

Here the divergence term can be estimated according to Lemma 5.5 and Lemma 5.6 as follows:

$$\begin{aligned} &\int_{\Sigma_2^t} \sum_{|I| \leq N} \text{div}_{\tilde{g}}(e^{-2\delta \log \rho_0} \mathcal{F}_{\tilde{g}}(T_2, \tilde{D}^I \tilde{h})) d\text{vol}_{\tilde{g}}^{T_2'} \\ &= \int_{\Sigma_2^t} \sum_{|I| \leq N} b^{-2\delta} (\mathcal{Q}_{\tilde{g}}(T_2, \tilde{D}^I \tilde{h}) - 2\delta \langle \mathcal{F}_{\tilde{g}}(T_2, \tilde{D}^I \tilde{h}), \nabla \log b \rangle_{\tilde{g}} + \langle \nabla T_2, \nabla \tilde{D}^I \tilde{h} \rangle_{\tilde{g}} (\square_{\tilde{g}} + \gamma_0) \tilde{D}^I \tilde{h}) d\text{vol}_{\tilde{g}}^{T_2'} \\ &\leq [a^{-1}(1 - 2\delta) + 2C'' a^{\delta - \delta' - 1} + 2C'' a^{\frac{n-4}{2} - \frac{3}{4} - \frac{1}{2}\delta'} M_2^N(\mathbf{t}; \tilde{h}, \tilde{g})] (M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \end{aligned}$$

where $C'' > 0$ is a constant independent of ϵ_2'' if it is small enough. Here $-\frac{3}{4} - \frac{1}{2}\delta' > \delta - \delta' - 1$ since $\delta - \delta' < \frac{1}{2} - \delta$. Notice the faster decaying of $\tilde{h}_{\rho\rho}$ and $\tilde{D} \tilde{h}_{\rho\rho}$ guarantee that the perturbation of $\mathcal{Q}_{\tilde{g}}, \mathcal{F}_{\tilde{g}}, \square_{\tilde{g}}$ all fall in the smaller term and the leading term $a^{-1}(1 - 2\delta)$ is determined by $\tilde{g} = \tilde{m}$. Here $\partial_t (L_2^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \geq 0$ since it is an increasing function of $\mathbf{t}$. Then

$$\begin{aligned} \partial_t M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) &\leq a^{-1}(\frac{1}{2} - \delta) M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) + C'' a^{\delta - \delta' - 1} M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) + C'' a^{-\frac{1}{2}} (M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \\ \implies \partial_t (a^{\frac{1}{2} - \delta} M_2^N(\mathbf{t}; \tilde{h}, \tilde{g})) &\leq C'' a^{\delta - \delta' - 1} (a^{\frac{1}{2} - \delta} M_2^N(\mathbf{t}; \tilde{h}, \tilde{g})) + C'' a^{\delta - 1} (a^{\frac{1}{2} - \delta} M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2. \end{aligned}$$

Assume $f(\mathbf{t}) = |\mathbf{t}|^{\frac{1}{2} - \delta} M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) < 1$, which is true at $\mathbf{t} = -\frac{1}{4}$ if ϵ_2'' small enough, then

$$\partial_t f \leq 2C'' |\mathbf{t}|^{\delta - \delta' - 1} f \implies f(\mathbf{t}) \leq e^{\frac{2C''}{\delta - \delta'} (\frac{1}{4})^{\delta - \delta'}} f(-\frac{1}{4}).$$

Hence $f(\mathbf{t}) < 1$ is true for all $\mathbf{t} \in [-\frac{1}{4}, \mathbf{t}_0)$ if ϵ_2'' small enough, which implies that

$$M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) \leq C'''' |\mathbf{t}|^{\delta - \frac{1}{2}} M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{g}), \quad \forall \mathbf{t} \in [-\frac{1}{4}, \mathbf{t}_0)$$

for some $C'''' > 0$ is independent of ϵ_2' if it is small enough. Notice that

$$\begin{aligned} \partial_{\mathbf{t}}(L_2^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 &\leq a^{2\delta-2}[(1-2\delta) + 2C'' a^{\delta-\delta'} + 2C'' C'''' a^{\delta} M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{g})](C'''' M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{g}))^2, \\ \implies L_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) &\leq C'''' |\mathbf{t}|^{\delta - \frac{1}{2}} M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{g}), \quad \forall \mathbf{t} \in [-\frac{1}{4}, \mathbf{t}_0) \end{aligned}$$

where $C'''' > 0$ is a constant independent of ϵ_2'' if it is small enough. Taking $C = C'''' + C'''''$, we prove the statement for $\mathbf{t} \in [-\frac{1}{4}, \mathbf{t}_0)$. Moreover,

$$\begin{aligned} \|\rho_0^{\frac{n-1}{2}} \tilde{h}\|_{L^\infty(\Sigma_2^1)} &\leq \|\tilde{h}\|_{L^\infty(\Sigma_1)} + C' \int_{-\frac{1}{4}}^{\mathbf{t}} (-T_2')^{-\frac{1}{2}} M_2^N(T_2'; \tilde{h}, \tilde{g}) dT_2' \leq \|\tilde{h}\|_{L^\infty(\Sigma_1)} + C C' (\frac{1}{4})^{\delta} \epsilon_2'', \\ \|\rho_0^{\frac{n-1}{2}} \rho_0^{-\delta} \tilde{h}_{\rho\rho}\|_{L^\infty(\bar{\Sigma}_2^1)} &\leq C' |\mathbf{t}|^{\frac{1}{2}-\delta} M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) \leq C C' \epsilon_2'', \end{aligned}$$

for some $C' > 0$ independent of ϵ_2'' if it is small enough. Hence choosing ϵ_2'' even smaller if necessary then Lemma 6.3 still holds at $T_2' = \mathbf{t}_0$, which allows us to extend the solution up to and hence a bit over $T_2' = \mathbf{t}_0$ with all the above estimates holds without changing of constants. Therefore the solution will never stop until it arrives $\tilde{\Omega}_2 \cap S_1$. We finish the proof. $\square$

From the the discussion in $\tilde{\Omega}_1$, we know that if the Cauchy data small enough, i.e. ϵ small enough, we have the condition in Proposition 6.2 is satisfied. Hence we can solve the equation (4.5) up to $\tilde{\Omega}_2 \cap S_1^+$.

Corollary 2. *For $n \geq 4$, suppose $M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{m}) < \epsilon_2''$ with ϵ_2'' chosen such that Proposition 6.2 holds. Then $\tilde{h}$ is $C^{0,\delta}$ up to $\tilde{\Omega}_2 \cap S_1^+$ and*

$$\|\tilde{h}\|_{S_1^+} \|\rho_1^{\delta} H_b^N(\tilde{\Omega}_2 \cap S_1^+)\| \leq C M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{m})$$

for some $C > 0$ is a constant depending on δ, δ' but independent of ϵ_2'' if it is small enough.

Proof. By Lemma 6.3 and Proposition 6.2, for some constant $C' > 0$

$$|\partial_a \tilde{h}| \leq C' a^{\delta-1} b^{\delta} M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{g}).$$

Hence $\tilde{h}$ is $C^{0,\delta}$ up to $\tilde{\Omega}_2 \cap S_1^+$ and the radiations field $\tilde{h}|_{\tilde{\Omega}_2 \cap S_1}$ is well defined. Moreover, define

$$Q_2^N(\mathbf{t}; v, \tilde{g}) = \left(\int_{\bar{\Sigma}_2^1} \sum_{k+|I| \leq N} |(b\partial_b)^k \partial_{\theta}^I v|^2 d\mu_{\tilde{g}}^{T_2'} \right)^{\frac{1}{2}}.$$

Then for some $C'' > 0$ and $C'''' > 0$

$$\partial_{\mathbf{t}}(Q_2^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \leq 2C'''' |\mathbf{t}|^{-\frac{1}{2}} M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) Q_2^N(\mathbf{t}; \tilde{h}, \tilde{g}), \quad Q_2^N(-\frac{1}{4}; \tilde{h}, \tilde{g}) \leq C'' M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{m}).$$

which implies that

$$Q_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) \leq \int_{-\frac{1}{4}}^0 C'''' |\mathbf{t}|^{-\frac{1}{2}} M_2^N(\mathbf{t}; \tilde{h}, \tilde{g}) + Q_2^N(-\frac{1}{4}; \tilde{h}, \tilde{g}) \leq C M_2^N(-\frac{1}{4}; \tilde{h}, \tilde{m}), \quad \forall \mathbf{t} \in [-\frac{1}{4}, 0].$$

Finally, $Q_2^N(0; \tilde{h}, \tilde{g})$ gives the norm bound of $\tilde{h}|_{S_1^+ \cap \tilde{\Omega}_2}$. $\square$

6.3. **In $\tilde{\Omega}_3$.** To solve the equations (4.5) in $\tilde{\Omega}_3$ we first notice that with Cauchy data small enough, the solution we find in $\tilde{\Omega}_1 \cup \tilde{\Omega}_2$ can be extended to $\tilde{\Omega}_3''$ since it is compact in the interior of X . We show that if the Cauchy data is small enough then this extension can be taken into $\tilde{\Omega}_3'$ and never stop until it arrives S_1^+ .

For $\mathfrak{t} \in [-1, 0]$, denote by

$$\tilde{\Sigma}_2^{\mathfrak{t}} = \Sigma_2 \cap \{-1 \leq T'_3 \leq \mathfrak{t}\}, \quad \Sigma_3^{\mathfrak{t}} = \{T'_3 = \mathfrak{t}\} \cap \tilde{\Omega}_3, \quad \Sigma_4^{\mathfrak{t}} = \Sigma_4 \cap \{-1 \leq T'_3 \leq \mathfrak{t}\}$$

the space like hypersurfaces w.r.t. $\tilde{m}$ with defining function T_2 , T'_3 and T_3 separately, and $\Omega_3^{\mathfrak{t}}$ the domain bounded by $\tilde{\Sigma}_2^{\mathfrak{t}}, \Sigma_3^{\mathfrak{t}}, \Sigma_4^{\mathfrak{t}}, \Sigma_3$. Define the high energy norm on $\Sigma_2^{\mathfrak{t}}, \Sigma_3^{\mathfrak{t}}, \Sigma_4^{\mathfrak{t}}$ by

$$\begin{aligned} M_3^N(\mathfrak{t}; v, \tilde{g}) &= \left(\int_{\Sigma_3^{\mathfrak{t}}} \sum_{|I| \leq N} \langle \mathcal{F}_{\tilde{g}}(T_3, \tilde{D}^I v), \nabla T'_3 \rangle_{\tilde{g}} d\mu_{\tilde{g}}^{T'_3} \right)^{\frac{1}{2}}, \\ L_3^N(\mathfrak{t}; v, \tilde{g}) &= \left(\int_{\Sigma_4^{\mathfrak{t}}} \sum_{|I| \leq N} \langle \mathcal{F}_{\tilde{g}}(T_3, \tilde{D}^I v), \nabla T_3 \rangle_{\tilde{g}} d\mu_{\tilde{g}}^{T_3} \right)^{\frac{1}{2}}, \\ \tilde{L}_3^N(\mathfrak{t}; v, \tilde{g}) &= \left(\int_{\Sigma_2^{\mathfrak{t}}} \sum_{|I| \leq N} \langle \mathcal{F}_{\tilde{g}}(T_3, \tilde{D}^I v), \nabla T_2 \rangle_{\tilde{g}} d\mu_{\tilde{g}}^{T_2} \right)^{\frac{1}{2}}, \end{aligned}$$

Here $d\mu_{\tilde{g}}^{T'_3} \wedge dT'_3 = d\text{vol}_{\tilde{g}}$, $d\mu_{\tilde{g}}^{T_3} \wedge dT_3 = d\text{vol}_{\tilde{g}}$, $d\mu_{\tilde{g}}^{T_2} \wedge dT_2 = d\text{vol}_{\tilde{g}}$ and $\tilde{D} \in \mathcal{B}_3$. Here

$$\mathcal{B}_3 = \{\partial_\tau, \rho\partial_\rho, Z_{ij} : i, j = 1, \dots, n\}.$$

Notice that the two quadratic forms $\langle \mathcal{F}_{\tilde{g}}(T_3, \tilde{D}^I v), \nabla T_2 \rangle_{\tilde{g}}$ and $\langle \mathcal{F}_{\tilde{g}}(T_2, \tilde{D}^I v), \nabla T_2 \rangle_{\tilde{g}}$ are equivalent on Σ_2 . We choose $\gamma_0 = -\frac{(n-1)(n-3)}{4}$ in the definition of $\mathcal{F}_{\tilde{g}}(T_3, v)$ in $\tilde{\Omega}_3'$. With above notation, by classical theorem for wave equations and Proposition 6.1 and Proposition 6.2, for all $\mathfrak{t} \in [-1, 0]$

$$(6.5) \quad \begin{aligned} M_3^N(-1; v, \tilde{m}) &< C_3\epsilon, \quad L_3^N(-1; v, \tilde{m}) < C_3\epsilon, \quad \tilde{L}_3^N(-1; v, \tilde{m}) = 0; \\ \partial_{\mathfrak{t}}(\tilde{L}_3^N(\mathfrak{t}; v, \tilde{g}))^2 &< C_3^2|\mathfrak{t}|^{2\delta-2}\epsilon^2, \quad |\tilde{h}_{\rho\rho}|_{\Sigma_2} + |\tilde{D}\tilde{h}_{\rho\rho}|_{\Sigma_2} \leq C_3\rho_1^\delta\epsilon, \end{aligned}$$

where $C_3 > 0$ independent of ϵ if it is small enough. In particular we choose $\delta' \in (0, \delta)$ in the definition of T_3 such that $\frac{1}{2} - \delta' < 1 - 2\delta$ and $\tau_0 > 8$ large in the definition of T'_3 . Similar as Lemma 6.3, we have

Lemma 6.4. *For $n \geq 4$ there exists $\epsilon_3 > 0$ such that if $\|\tilde{h}\|_{L^\infty(\Sigma_3^{\mathfrak{t}})} + \|\rho_1^{-\delta}\tilde{h}_{\rho\rho}\|_{L^\infty(\Sigma_3^{\mathfrak{t}})} < \epsilon_3$, then*

$$\begin{aligned} \frac{1}{\sqrt{2}}M_3^N(\mathfrak{t}; v, \tilde{m}) &\leq M_3^N(\mathfrak{t}; v, \tilde{g}) \leq \sqrt{2}M_3^N(\mathfrak{t}; v, \tilde{m}), \\ \sum_{|I| \leq \frac{N}{2}+1} |\tilde{D}^I v| &\leq CM_3^N(\mathfrak{t}; v, \tilde{g}), \quad \sum_{|I| \leq \frac{N}{2}} |\tilde{D}^I \partial_a v| \leq C\rho_1^{-\frac{1}{2}}M_3^N(\mathfrak{t}; v, \tilde{g}), \\ |\tilde{h}_{\rho\rho}| + |\tilde{D}\tilde{h}_{\rho\rho}| &\leq C_3\rho_1^\delta\epsilon + C\rho_1^{\frac{1}{2}}M_3^N(\mathfrak{t}; \tilde{h}, \tilde{m}). \end{aligned}$$

for some constant C independent of ϵ_3 if it is small enough.

Proposition 6.3. *For $n \geq 4$, there exists $\epsilon_3'' > 0$ such that if $\epsilon \leq \epsilon_3''$, then the equations (4.5) have a unique solution up to Σ_4 such that*

$$M_3^N(\mathfrak{t}; \tilde{h}, \tilde{g}) + L_3^N(\mathfrak{t}; \tilde{h}, \tilde{g}) \leq C|\mathfrak{t}|^{\delta-\frac{1}{2}}\epsilon, \quad \forall \mathfrak{t} \in [-1, 0],$$

where $C > 0$ is a constant depending on δ, δ' but independent of ϵ_3'' if it is small enough.

Proof. We first can choose ϵ_3'' such that we can solve the equation in $\tilde{\Omega}_1 \cup \tilde{\Omega}_2 \cup \tilde{\Omega}_3''$ and (6.5) holds. Choose ϵ_3'' even smaller if necessary such that we can extend the solution to $\Sigma_3^{\mathfrak{t}_0}$ for some $\mathfrak{t}_0 \in (-1, 0)$ with Lemma 6.4 satisfied and

$$|\mathfrak{t}|^{\frac{1}{2}-\delta}M_3^N(\mathfrak{t}; \tilde{h}, \tilde{g}) < 1.$$

Then By Lemma 5.2,

$$\begin{aligned} & (M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 + (L_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 - (L_3^N(-1; \tilde{h}, \tilde{g}))^2 - (M_3^N(-1; \tilde{h}, \tilde{g}))^2 - (\tilde{L}_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \\ &= \int_{\Omega_3^t} \sum_{|I| \leq N} \operatorname{div}_{\tilde{g}}(\mathcal{F}_{\tilde{g}}(T_3, \tilde{D}^I \tilde{h})) d\operatorname{vol}_{\tilde{g}}. \end{aligned}$$

Here the divergence term can be estimated by Lemma 5.7 and Lemma 5.8 as follows:

$$\begin{aligned} & \int_{\Sigma_3^t} \sum_{|I| \leq N} \operatorname{div}_{\tilde{g}}(\mathcal{F}_{\tilde{g}}(T_3, \tilde{D}^I \tilde{h})) d\operatorname{vol}_{\tilde{g}}^{T'_3} \\ &= \int_{\Sigma_3^t} \sum_{|I| \leq N} (\mathcal{Q}_{\tilde{g}}(T_3, \tilde{D}^I \tilde{h}) + \langle \nabla T_3, \nabla \tilde{D}^I \tilde{h} \rangle_{\tilde{g}} (\square_{\tilde{g}} + \gamma_0) \tilde{D}^I \tilde{h}) d\operatorname{vol}_{\tilde{g}}^{T'_3} \\ &\leq [(\frac{1}{2} - \delta') |\mathbf{t}|^{-1} + 2C' |\mathbf{t}|^{\delta - \delta' - 1} + 2C' |\mathbf{t}|^{-\frac{1}{2}} M_3^N(\mathbf{t}; \tilde{h}, \tilde{g})] (M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \\ &\leq [(\frac{1}{2} - \delta') |\mathbf{t}|^{-1} + 4C' |\mathbf{t}|^{\delta - \delta' - 1}] (M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2. \end{aligned}$$

Hence

$$\begin{aligned} & \partial_t (M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \leq C_3^2 |\mathbf{t}|^{2\delta - 2} \epsilon^2 + [(\frac{1}{2} - \delta') |\mathbf{t}|^{-1} + 4C' |\mathbf{t}|^{\delta - \delta' - 1}] (M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \\ \implies & \partial_t (e^{\frac{4C'}{\delta - \delta'} \rho^{\delta - \delta'}} |\mathbf{t}|^{\frac{1}{2} - \delta'} (M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2) \leq C_3^2 \epsilon^2 |\mathbf{t}|^{2\delta - \delta' - \frac{3}{2}} \\ \implies & (M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \leq \frac{C_3^2 \epsilon^2}{(\frac{1}{2} - \delta') - (1 - 2\delta)} |\mathbf{t}|^{2\delta - 1} + e^{\frac{4C'}{\delta - \delta'}} |\mathbf{t}|^{\delta' - \frac{1}{2}} (M_3^N(-1; \tilde{h}, \tilde{g}))^2 \leq C_1^2 \epsilon^2 |\mathbf{t}|^{2\delta - 1}. \end{aligned}$$

Here $\frac{1}{2} - \delta' \in (0, 1 - 2\delta)$ and C depends on δ, δ' only if ϵ_3'' small enough. Notice that, by Lemma 6.4,

$$\begin{aligned} & \|\tilde{h}\|_{L^\infty(\Sigma_3^t)} \leq \|\tilde{h}\|_{L^\infty(\Sigma_3^{-1})} + C'' \int_{-1}^t |T_3'|^{-\frac{1}{2}} M_3^N(T_3'; \tilde{h}, \tilde{g}) dT_3' \leq \|\tilde{h}\|_{L^\infty(\Sigma_3^{-1})} + \frac{1}{\delta} C'' C_1 \epsilon, \\ & \|\rho_1^{-\delta} \tilde{h}_{\rho\rho}\|_{L^\infty(\Sigma_3^t)} \leq \|\rho_1^{-\delta} \tilde{h}_{\rho\rho}\|_{L^\infty(\tilde{\Sigma}_2^t)} + C'' |\mathbf{t}|^{\frac{1}{2} - \delta} M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}) \leq \|\rho_1^{-\delta} \tilde{h}_{\rho\rho}\|_{L^\infty(\tilde{\Sigma}_2^t)} + C_1 C'' \epsilon. \end{aligned}$$

Choose ϵ_3'' even smaller if necessary such that $M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}) < C_1 \epsilon |\mathbf{t}|^{\delta - \frac{1}{2}}$ implies Lemma 6.4. Then when we extend the solution over $\mathbf{t}_0$, all the above inequalities still hold with the same constants and hence the extension does not stop until it arrives S_1^+ . Moreover,

$$\begin{aligned} & \partial_t ((L_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2) \leq [(\frac{1}{2} - \delta') |\mathbf{t}|^{-1} + 4C' |\mathbf{t}|^{\delta - \delta' - 1}] (M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \leq C''' \epsilon^2 |\mathbf{t}|^{2\delta - 2}, \\ \implies & L_3^N(\mathbf{t}; \tilde{h}, \tilde{g}) \leq C_2 \epsilon |\mathbf{t}|^{\delta - \frac{1}{2}}. \end{aligned}$$

Let $C = C_1 + C_2$. We finish the proof. $\square$

Corollary 3. For $n \geq 4$ suppose $\epsilon < \epsilon_3''$ with ϵ_3'' chosen in Proposition 6.3. Then $\tilde{h}$ is C^δ up to $\tilde{\Omega}_3 \cap S_1^+$ and

$$\|\tilde{h}_{S_1^+}\|_{H^N(\tilde{\Omega}_3 \cap S_1^+)} \leq C\epsilon$$

for some constant $C > 0$ depending on δ but independent of ϵ_3'' if it is small enough.

Proof. By Lemma 6.4 and Proposition 6.3, for some constant $C' > 0$

$$|\partial_\rho \tilde{h}| \leq C' \epsilon \rho^{\delta - 1}.$$

Hence $\tilde{h}$ is $C^{0, \delta}$ up to $\tilde{\Omega}_3 \cap S_1^+$ and the radiations field $\tilde{h}|_{\tilde{\Omega}_3 \cap S_1}$ is well defined. Moreover, define

$$Q_3^N(\mathbf{t}; v, \tilde{g}) = \left(\int_{\Sigma_3^t} \sum_{k+|I| \leq N} |\partial_\tau^k \partial_\theta^I v|^2 d\mu_{\tilde{g}}^{T'_3} \right)^{\frac{1}{2}}.$$

Then for some $C'' > 0$ and $C''' > 0$

$$\partial_t (Q_3^N(\mathbf{t}; \tilde{h}, \tilde{g}))^2 \leq 2C''' |\mathbf{t}|^{-\frac{1}{2}} M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}) Q_3^N(\mathbf{t}; \tilde{h}, \tilde{g}), \quad Q_3^N(-1; \tilde{h}, \tilde{g}) \leq C'' M_3^N(-1; \tilde{h}, \tilde{m}).$$

which implies that

$$Q_3^N(\mathbf{t}; \tilde{h}, \tilde{g}) \leq \int_{-1}^0 C''' |\mathbf{t}|^{-\frac{1}{2}} M_3^N(\mathbf{t}; \tilde{h}, \tilde{g}) + Q_3^N(-1; \tilde{h}, \tilde{g}) \leq C\epsilon, \quad \forall \mathbf{t} \in [-1, 0].$$

Finally, $Q_3^N(0; \tilde{h}, \tilde{g})$ gives the norm bound of $\tilde{h}|_{S_1^+ \cap \tilde{\Omega}_3}$. $\square$

6.4. In $\tilde{\Omega}_4$. To Solve the equation (4.5) in $\tilde{\Omega}_4$, we first notice that if the Cauchy data is small enough, then we can extend the solution a bit over Σ_4 on a compact region. We show that if $\epsilon > 0$ small enough, then this extension can be taken all over $\tilde{\Omega}_4$ and hence we solve the equation globally.

Let $\mathfrak{t}_0 = \inf\{\mathfrak{t} : \tilde{\Sigma}_4 \cap \tilde{\Omega}_4 \neq \emptyset\}$. For $\mathfrak{t} \in [\mathfrak{t}_0, 0]$, denote by

$$\tilde{\Sigma}_4^{\mathfrak{t}} = \Sigma_4 \cap \{-1 \leq T'_4 \leq \mathfrak{t}\}, \quad \tilde{\Sigma}_4^{\mathfrak{t}} = \tilde{\Omega}_4 \cap \{T'_4 = \mathfrak{t}\}$$

the space-like hypersurfaces w.r.t $\tilde{m}$ with defining function T_3 and T'_4 separately, and $\Omega_4^{\mathfrak{t}}$ the domain bounded by $\tilde{\Sigma}_4^{\mathfrak{t}}$ and $\tilde{\Sigma}_4^{\mathfrak{t}}$. Define the high energy norm on $\tilde{\Sigma}_4^{\mathfrak{t}}$ and $\tilde{\Sigma}_4^{\mathfrak{t}}$ by

$$M_4^N(\mathfrak{t}; v, \tilde{g}) = \left(\int_{\tilde{\Sigma}_4^{\mathfrak{t}}} \sum_{|I| \leq N} \langle \mathcal{F}_{\tilde{g}}(T_4, \tilde{D}^I v), \nabla T'_4 \rangle_{\tilde{g}} d\mu_{\tilde{g}}^{T'_4} \right)^{\frac{1}{2}},$$

$$L_4^N(\mathfrak{t}; v, \tilde{g}) = \left(\int_{\tilde{\Sigma}_4^{\mathfrak{t}}} \sum_{|I| \leq N} \langle \mathcal{F}_{\tilde{g}}(T_4, \tilde{D}^I v), \nabla T_3 \rangle_{\tilde{g}} d\mu_{\tilde{g}}^{T_3} \right)^{\frac{1}{2}},$$

Here $d\mu_{\tilde{g}}^{T'_4} \wedge dT'_4 = d\text{vol}_{\tilde{g}}$, $d\mu_{\tilde{g}}^{T_3} \wedge dT_3 = d\text{vol}_{\tilde{g}}$ and $\tilde{D} \in \mathcal{B}_4 = \{Z_{\mu\nu} : \mu, \nu = 0, \dots, n\}$. Notice that the two quadratic forms $\langle \mathcal{F}_{\tilde{g}}(T_3, \tilde{D}^I v), \nabla T_3 \rangle_{\tilde{g}}$ and $\langle \mathcal{F}_{\tilde{g}}(T_4, \tilde{D}^I v), \nabla T_3 \rangle_{\tilde{g}}$ are equivalent on Σ_4 . We choose $\gamma_0 = -\frac{n^2-1}{4}$ in the definition of $\mathcal{F}_{\tilde{g}}(T_4, v)$ in $\tilde{\Omega}_4$. With above notation, by Proposition 6.1, 6.2, 6.3, for all $\mathfrak{t} \in [\mathfrak{t}_0, 0]$

$$(6.6) \quad M_4^N(\mathfrak{t}_0; v, \tilde{g}) = 0, \quad L_4^N(\mathfrak{t}_0; v, \tilde{g}) = 0,$$

$$L_4^N(\mathfrak{t}; v, \tilde{g}) < C_4 |\mathfrak{t}|^{\delta - \frac{1}{2}} \epsilon, \quad |\tilde{h}_{\rho\rho}|_{\Sigma_4} + |\tilde{D}\tilde{h}_{\rho\rho}|_{\Sigma_4} \leq C_4 \rho_1^\delta \epsilon$$

where $C_4 > 0$ independent of ϵ if it is small enough. In particular, in the definition of T_4 we choose $\delta' \in (0, \delta)$ such that $\delta - \delta' < \frac{1}{2} - \delta$ and in the definition of T'_4 we choose $\alpha \in (1, \frac{n-1}{1-2\delta})$ such that Lemma 5.10 holds for $\lambda = 1 - 2\delta$. Similar as Lemma 6.4, we have

Lemma 6.5. *For $n \geq 4$ there exists $\epsilon_4 > 0$ such that if $\|\rho_2^{\frac{n-1}{2}} \tilde{h}\|_{L^\infty(\tilde{\Sigma}_4^{\mathfrak{t}})} + \|\rho_1^{-\delta} \rho_2^{\frac{n-1}{2}} \tilde{h}_{\rho\rho}\|_{L^\infty(\tilde{\Sigma}_4^{\mathfrak{t}})} < \epsilon_4$, then*

$$\frac{1}{\sqrt{2}} M_4^N(\mathfrak{t}; v, \tilde{m}) \leq M_4^N(\mathfrak{t}; v, \tilde{g}) \leq \sqrt{2} M_4^N(\mathfrak{t}; v, \tilde{m}),$$

$$\sum_{|I| \leq \frac{N}{2} + 1} |\tilde{D}^I v| \leq C M_4^N(\mathfrak{t}; v, \tilde{g}), \quad \sum_{|I| \leq \frac{N}{2}} |\tilde{D}^I \partial_a v| \leq C \rho_1^{-\frac{1}{2}} M_4^N(\mathfrak{t}; v, \tilde{g}),$$

$$|\tilde{h}_{\rho\rho}| + |\tilde{D}\tilde{h}_{\rho\rho}| \leq C_4 \rho_1^\delta \epsilon + C \rho_1^{\frac{1}{2}} (\ln \rho_2) M_4^N(\mathfrak{t}; \tilde{h}, \tilde{m}).$$

for some constant C independent of ϵ_4 if it is small enough.

Proposition 6.4. *For $n \geq 4$, there exists $\epsilon_4'' > 0$ such that if $\epsilon \leq \epsilon_4''$, then the equations (4.5) have a unique solution all over $\tilde{\Omega}_4$ such that*

$$M_4^N(\mathfrak{t}; \tilde{h}, \tilde{g}) \leq C |\mathfrak{t}|^{\delta - \frac{1}{2}} \epsilon, \quad \forall \mathfrak{t} \in [\mathfrak{t}_0, 0],$$

where $C > 0$ is a constant depending on $\delta - \delta'$ but independent of ϵ_4'' if it is small enough.

Proof. We first can choose ϵ_4'' such that we can solve the equation in $\tilde{\Omega}_1 \cup \tilde{\Omega}_2 \cup \tilde{\Omega}_3$ and (6.6) holds. Choose ϵ_4'' even smaller if necessary such that we can extend the solution to $\tilde{\Sigma}_4^{\mathfrak{t}_1}$ for some $\mathfrak{t}_1 \in (\mathfrak{t}_0, 0)$ with Lemma 6.5 satisfied and

$$|\mathfrak{t}|^{\frac{1}{2} - \delta} M_4^N(\mathfrak{t}; \tilde{h}, \tilde{g}) < 1.$$

Then By Lemma 5.2,

$$(M_4^N(\mathfrak{t}; \tilde{h}, \tilde{g}))^2 - (L_4^N(\mathfrak{t}; \tilde{h}, \tilde{g}))^2 = \int_{\Omega_4^{\mathfrak{t}}} \sum_{|I| \leq N} \text{div}_{\tilde{g}}(\mathcal{F}_{\tilde{g}}(T_4, \tilde{D}^I \tilde{h})) d\text{vol}_{\tilde{g}}.$$

Here the divergence term can be estimated by Lemma 5.9, Lemma 5.10 and Lemma 5.11 as follows:

$$\begin{aligned}
& \int_{\tilde{\Sigma}_4^t} \sum_{|I| \leq N} \operatorname{div}_{\tilde{g}}(F(T_4, \tilde{D}^I \tilde{h})) d\operatorname{vol}_{\tilde{g}}^{T_4'} \\
&= \int_{\tilde{\Sigma}_4^t} \sum_{|I| \leq N} (\mathcal{Q}_{\tilde{g}}(T_4, \tilde{D}^I \tilde{h}) + \langle \nabla T_4, \nabla \tilde{D}^I \tilde{h} \rangle_{\tilde{g}} (\square_{\tilde{g}} + \gamma_0) \tilde{D}^I \tilde{h}) d\operatorname{vol}_{\tilde{g}}^{T_4'} \\
&\leq [(1-2\delta)|t|^{-1} + 2C'|t|^{\delta-\delta'-1} + 2C'|t|^{-\frac{1}{2}} M_4^N(t; \tilde{h}, \tilde{g})] (M_4^N(t; \tilde{h}, \tilde{g}))^2 \\
&\leq [(1-2\delta)|t|^{-1} + 4C'|t|^{\delta-\delta'-1}] (M_4^N(t; \tilde{h}, \tilde{g}))^2.
\end{aligned}$$

Hence

$$\begin{aligned}
& \partial_t (M_4^N(t; \tilde{h}, \tilde{g}))^2 \leq \partial_t (L_4^N(t; \tilde{h}, \tilde{g}))^2 + [(1-2\delta)|t|^{-1} + 4C'|t|^{\delta-\delta'-1}] (M_4^N(t; \tilde{h}, \tilde{g}))^2 \\
\Rightarrow & \partial_t (e^{\frac{4C'}{\delta-\delta'}} |t|^{\delta-\delta'} |t|^{1-2\delta} (M_4^N(t; \tilde{h}, \tilde{g}))^2) \leq e^{\frac{4C'}{\delta-\delta'}} |t|^{\delta-\delta'} |t|^{1-2\delta} \partial_t (L_4^N(t; \tilde{h}, \tilde{g}))^2 \\
\Rightarrow & M_4^N(t; \tilde{h}, \tilde{g}) \leq e^{\frac{2C'}{\delta-\delta'} |t_0|^{\delta-\delta'}} L_4^N(t; \tilde{h}, \tilde{g}) \leq C |t|^{\delta-\frac{1}{2}} \epsilon.
\end{aligned}$$

Here C depends on δ, δ' only if ϵ_4'' small enough. By Lemma 6.5, this implies

$$|\rho_2^{\frac{n-1}{2}} \tilde{h}| \leq C'' \rho_2^{\sigma'} \epsilon, \quad |\rho_2^{\frac{n-1}{2}} \tilde{\partial} \tilde{h}| \leq C'' \rho_2^{\sigma'} \epsilon, \quad \text{and} \quad |\rho_2^{\frac{n-1}{2}} \tilde{h}_{\rho\rho}| < C'' \rho_1^\delta \rho_2^{\sigma'} (\ln \rho_2) \epsilon,$$

where $\sigma' = \frac{n-1}{2} - \alpha(\frac{1}{2} - \delta) > 0$, and C'' only depend on δ, δ', α if ϵ is small enough. Choose ϵ_4'' even smaller if necessary such that $M_4^N(t; \tilde{h}, \tilde{g}) < C\epsilon |t|^{\delta-\frac{1}{2}}$ implies Lemma 6.5. Then when we extend the solution over t_1 , all the above inequalities still hold with same constants and hence the extension does not stop. We solve the equation (4.5) globally. $\square$

Corollary 4. *For $n \geq 4$ suppose $\epsilon < \epsilon_4''$ with ϵ_4'' chosen in Proposition 6.4. Then $\tilde{h}$ is C^δ up to $\tilde{\Omega}_4 \cap S_1^+$ and for $0 < \sigma < \frac{n-1}{2} - \alpha(\frac{1}{2} - \delta)$,*

$$\|\tilde{h}_{S_1^+}\|_{\rho_2^{\frac{n-1}{2}-\sigma} H_b^N(\tilde{\Omega}_4 \cap S_1^+)} \leq C\epsilon$$

for some constant $C > 0$ depending on $\alpha, \delta, \delta', \sigma$ but independent of ϵ_4'' if it is small enough.

Proof. By Lemma 6.5 and Proposition 6.4, in $\tilde{\Omega}_4$ for some constant $C' > 0$

$$\sum_{|I| \leq \frac{N}{2}} |\tilde{D}^I \tilde{h}| \leq C' \bar{a}^{\delta-\frac{1}{2}} \bar{b}^{\alpha(\delta-\frac{1}{2})} \epsilon, \quad \sum_{|I| \leq \frac{N}{2}} |\partial_{\bar{a}} \tilde{D}^I \tilde{h}| \leq C' \bar{a}^{\delta-1} \bar{b}^{\alpha(\delta-\frac{1}{2})} \epsilon.$$

Hence $\tilde{h}$ is $C^{0,\delta}$ up to $\tilde{\Omega}_4 \cap S_1^+$ and the radiations field $\tilde{h}|_{\tilde{\Omega}_4 \cap S_1}$ is well defined. Moreover, define

$$Q_4^N(t; v, \tilde{g}) = \left(\int_{\tilde{\Sigma}_4^t} \bar{b}^{n-1-2\sigma+\alpha} \sum_{k+|I| \leq N} |(\bar{b}\partial_{\bar{b}})^k (\frac{\partial_{\bar{b}}}{1-\bar{a}})^I v|^2 d\mu_{\tilde{g}}^{T_4'} \right)^{\frac{1}{2}}.$$

Then for some $C'' > 0$

$$\partial_t (Q_4^N(t; \tilde{h}, \tilde{g}))^2 \leq 2C'' |t|^{\frac{n-1-2\sigma}{2\alpha}-1} M_2^N(t; \tilde{h}, \tilde{g}) Q_4^N(t; v, \tilde{g}), \quad Q_4^N(t_0; \tilde{h}, \tilde{g}) = 0.$$

Here $\frac{n-1-2\sigma}{2\alpha} > \frac{1}{2} - \delta$, which implies that

$$Q_4^N(t; \tilde{h}, \tilde{g}) \leq \int_{t_0}^0 C'' |t|^{\frac{n-1-2\sigma}{2\alpha}-1} M_4^N(t; \tilde{h}, \tilde{g}) \leq C\epsilon, \quad \forall t \in [t_0, 0],$$

where C depend on $\alpha, \delta, \delta', \sigma$. Finally, $Q_4^N(0; \tilde{h}, \tilde{g})$ gives the norm bound of $\tilde{h}|_{S_1^+ \cap \tilde{\Omega}_4}$. $\square$

7. NONLINEAR MØLLER WAVE OPERATOR

For $n \geq 4$, suppose $(h^0, h^1) \in \tilde{\mathcal{V}}_\epsilon^{N,\delta}$ with $N \geq n+6$, $\delta \in (0, \frac{1}{2})$ and $\epsilon > 0$ small such that Theorem 3 holds. Then the global solution h to reduced Einstein equation (1.3) obtained in Theorem 3 provides a true solution to Einstein equation (1.1) since h satisfies the harmonic gauge condition (1.4) globally by Theorem 1. This imposes constraint conditions on the radiation field.

$$\partial_\tau(\tilde{h}_{\mu\rho}|_{S_1^+}) - \frac{1}{2}\theta_\mu\partial_\tau(\text{tr}_m\tilde{h}|_{S_1^+}) = 0, \quad \mu = 0, 1, \dots, n.$$

This is equivalent to

$$(7.1) \quad (\tilde{h}_{\mu\rho}|_{S_1^+}) - \frac{1}{2}\theta_\mu(\text{tr}_m\tilde{h}|_{S_1^+}) = 0, \quad \mu = 0, 1, \dots, n.$$

when $\tilde{h}|_{S_1^+}$ has some decay as $|\tau| \rightarrow 0$. To refine the image space for the radiation field, first let us consider the Cauchy data with only conormal singularity at $\partial\Sigma_0$.

Lemma 7.1. *If $(h_0, h_1) \in \tilde{\mathcal{V}}_\epsilon^{N,\delta} \cap \mathcal{A}^{\frac{n-1}{2}+\delta}(\Sigma_0)$, then $h \in \mathcal{A}^{\frac{n-1}{2}+\delta, \frac{n-1}{2}, \sigma}(X)$, where $\sigma > 0$ is given in Theorem 3.*

Proof. For all $N' > N$, we estimate $M_i^{N'}(\mathbf{t}; \tilde{h}, \tilde{g})$ in the same way as we do for $M_i^N(\mathbf{t}; \tilde{h}, \tilde{g})$. The only difference is that in this case, we can replace the cubic term $(M_i^N(\mathbf{t}; \tilde{h}, \tilde{g}))^s$ in the estimates of divergence term by $C'M_i^N(\mathbf{t}, \tilde{h})(M_i^{N'}(\mathbf{t}, \tilde{h}))^2$ as a result of interpolation. Hence the nonlinear estimating reduces to linear estimating:

$$\begin{aligned} \partial_t M_1^{N'}(T_1; \tilde{h}, \tilde{g}) &\leq C' M_1^{N'}(T_1; \tilde{h}, \tilde{g}), \\ \partial_t M_i^{N'}(T_i; \tilde{h}, \tilde{g}) &\leq ((\tfrac{1}{2} - \delta)|T_i'|^{-1} + C'|T_i'|^{\delta-\delta'-1}) M_i^{N'}(T_i'; \tilde{h}, \tilde{g}), \quad i = 2, 3, 4 \end{aligned}$$

for some $C' > 0$. For some $C_1, C_2, C_3, C_4 > 0$, we have

$$\begin{aligned} M_1^{N'}(T_1; \tilde{h}, \tilde{g}) &\leq C_1 M_1^{N'}(0; \tilde{h}, \tilde{g}), \\ M_i^{N'}(T_i; \tilde{h}, \tilde{g}) &\leq C_i |T_i'|^{\delta-\frac{1}{2}} M_1^{N'}(0; \tilde{h}, \tilde{g}), \quad i = 2, 3, 4. \end{aligned}$$

Then by the same proof as the proof of Corollary 2, 3, 4, we have

$$\|\tilde{h}_{S_1^+}\|_{\rho_0^\delta \rho_2^{\sigma-\frac{n-1}{2}} H_b^{N'}(S_1^+)} \leq C M_1^{N'}(0; \tilde{h}, \tilde{g})$$

for some C only depending on N' . Since N' is arbitrary large, $\tilde{h}$ only have conormal singularity at ∂X . By Lemma 2.2, we have $\tilde{h} \in \mathcal{A}^{\delta, 0, \sigma-\frac{n-1}{2}}$. Hence $h = \tilde{\rho}^{\frac{n-1}{2}} \tilde{h} \in \mathcal{A}^{\frac{n-1}{2}+\delta, \frac{n-1}{2}, \sigma}(X)$. $\square$

Lemma 7.2. *If $(h_0, h_1) \in \tilde{\mathcal{V}}_\epsilon^{N,\delta} \cap \mathcal{A}^{\frac{n-1}{2}+\delta}(\Sigma_0)$, then $h \in \mathcal{A}^{\frac{n-1}{2}+\delta, \frac{n-1}{2}, \frac{n-1}{2}+\delta}(X)$.*

Proof. Applying the linear wave operator $\square_m$ to the solution h gives

$$\square_m h_{\mu\nu} = \square_g h_{\mu\nu} + (\square_m - \square_g) h_{\mu\nu} = F_{\mu\nu}(\partial h, \partial h) - H^{\alpha\beta} \partial_\alpha \partial_\beta h_{\mu\nu} \in \mathcal{A}^{n+1+2\delta, n-1, 2+2\sigma}(X).$$

Hence $h \in \mathcal{A}^{\frac{n-1}{2}+\delta, \frac{n-1}{2}, 2\delta}(X)$ if $2\sigma < \frac{n-1}{2} + \delta$. See [MW] for details. Repeating k times until $2^k \sigma \geq \frac{n-1}{2} + \delta$ and finally we have $h \in \mathcal{A}^{\frac{n-1}{2}+\delta, \frac{n-1}{2}, \frac{n-1}{2}+\delta}(X)$ $\square$

Fix Cauchy data $(h^0, h^1) \in \tilde{\mathcal{V}}_\epsilon^{N,\delta} \cap \mathcal{A}^{\frac{n-1}{2}+\delta}(\Sigma_0)$ and let h be the solution obtained in Theorem 3. Consider the Cauchy problem for linear wave equation with background metric $g = m + h$:

$$(7.2) \quad \square_g k_{\mu\nu} = F_{\mu\nu}(h)(\partial k, \partial h), \quad (k_{t=0}, \partial_t k|_{t=0}) = (k^0, k^1).$$

Then k globally exists and the radiation field is well defined. Moreover, $k = h$ is a solution if $(k^0, k^1) = (h^0, h^1)$. Denote by $\tilde{k} = \tilde{\rho}^{\frac{1-n}{2}} k$ and define the following map

$${}^g\mathcal{R}_{\mathcal{F}} : \rho_0^\lambda C^\infty \times \rho_0^{\lambda+1} C^\infty \ni (k^0, k^1) \rightarrow \tilde{k}|_{S_1} \in (\rho_0 \rho_2)^{\lambda-\frac{n-1}{2}} C^\infty(S_1^+).$$

for $\lambda \in \mathbb{C}$, $\Re \lambda \in (\frac{n-1}{2}, \frac{n}{2})$.

Lemma 7.3. *For $\Re \lambda \in (\frac{n-1}{2}, \frac{n+1}{2})$, ${}^g\mathcal{R}_{\mathcal{F}}$ and ${}^m\mathcal{R}_{\mathcal{F}}$ have the same boundary operator.*

Proof. With $(k^0, k^1) \in \rho_0^\lambda C^\infty \times \rho_0^{\lambda+1} C^\infty$, by similar energy estimates in the proof of Lemma 7.1, we first have $k \in \mathcal{A}^{\lambda, \frac{n-1}{2}, \sigma}$. Then applying $\square_m$ as in the proof of Lemma 7.2, we get $k \in \mathcal{A}^{\lambda, \frac{n-1}{2}, \lambda}$. Consider the equation

$$\square_m k'_{\mu\nu} = 0, \quad (k'_{t=0}, \partial_t k'|_{t=0}) = (k^0, k^1).$$

Then

$$\square_m(k - k') = -H^{\mu\nu} \partial_\mu \partial_\nu k + F(h)(\partial k, \partial h) \in \mathcal{A}^{\frac{n-1}{2}+\delta+\tau+2, n-1, \frac{n-1}{2}+\delta+\lambda+2}.$$

Hence $k' - k \in \mathcal{A}^{\frac{n-1}{2}+\delta+\lambda, \frac{n-1}{2}, \frac{n-1}{2}+\delta+\lambda} + \mathcal{A}^{\frac{n-1}{2}+\delta+\lambda+2, \frac{n-1}{2}, n-2}$. Since $\Re \lambda < n - 2$, this term contributes zero to the boundary operator. See [MW] for details. $\square$

By Lemma 7.3 and a similar proof for the mapping property of ${}^m\mathcal{R}_{\mathcal{F}}$ in [MW], we show the following lemma.

Lemma 7.4. *The linear Møller wave operator ${}^g\mathcal{R}_{\mathcal{F}}$ defines a continuous map for $\delta \in (0, \frac{1}{2})$*

$$\begin{aligned} {}^g\mathcal{R}_{\mathcal{F}} : \rho_0^{\frac{n-1}{2}+\delta} H_b^{N+1}(\Sigma_0) \times \rho_0^{\frac{n+1}{2}+\delta} H_b^{N+1}(\Sigma_0) \\ \longrightarrow (\rho_0 \rho_2)^\delta [H_b^{\frac{1}{2}-\delta}(\overline{\mathbb{R}}; H^{N+\frac{1}{2}+\delta}(\mathbb{S}^{n-1})) \cap L^2(\mathbb{S}^{n-1}; H_b^{N+1}(\overline{\mathbb{R}}))] \end{aligned}$$

Here $\|{}^g\mathcal{R}_{\mathcal{F}}\| < C$ for some constant $C > 0$ independent of $(h^0, h^1) \in \widetilde{\mathcal{V}}_\epsilon^{N, \delta} \cap \mathcal{A}^{\frac{n-1}{2}+\delta}(\Sigma_0)$. Moreover, ${}^m\mathcal{R}_{\mathcal{F}}$ is an isomorphism.

Denote by $\widetilde{\mathcal{W}}_\epsilon^{N, \delta}$ be the collection of elements in

$$(\rho_0 \rho_2)^\delta [H_b^{\frac{1}{2}-\delta}(\overline{\mathbb{R}}; H^{N+\frac{1}{2}+\delta}(\mathbb{S}^{n-1})) \cap L^2(\mathbb{S}^{n-1}; H_b^{N+1}(\overline{\mathbb{R}}))]$$

with norm less than ϵ and satisfying the harmonic gauge condition (7.1). Since $\delta > 0$, $\widetilde{\mathcal{W}}_\epsilon^{N, \delta}$ is a small neighborhood of 0 in the Sobolev space $\widetilde{\mathcal{W}}_\infty^{N, \delta}$.

Theorem 4. *With the assumption in Theorem 3 and choosing $\epsilon > 0$ even smaller if necessary, the nonlinear Møller wave operator defines a continuous map and open map*

$$\mathcal{R}_{\mathcal{F}} : \widetilde{\mathcal{V}}_\epsilon^{N, \delta} \longrightarrow \widetilde{\mathcal{W}}_{C\epsilon}^{N, \delta}$$

for some $C > 0$.

Proof. First, $\mathcal{R}_{\mathcal{F}}$ is well defined and continuous. This is obvious for $(h^0, h^1) \in \widetilde{\mathcal{V}}_\epsilon^{N, \delta} \cap \mathcal{A}^{\frac{n-1}{2}+\delta}(\Sigma_0)$ since

$$\mathcal{R}_{\mathcal{F}}(h^0, h^1) = {}^g\mathcal{R}_{\mathcal{F}}(h^0, h^1) < C\epsilon.$$

where C is given in Lemma 7.4. By density argument, it holds for all $(h^0, h^1) \in \widetilde{\mathcal{V}}_\epsilon^{N, \delta}$. Secondly, the linearization of $\mathcal{R}_{\mathcal{F}}$ at $(0, 0)$ is ${}^m\mathcal{R}_{\mathcal{F}}|_{\mathcal{T}^{N, \frac{n-1}{2}+\delta}}$, where $\mathcal{T}^{N, \frac{n-1}{2}+\delta}$ is the tangent space of $\widetilde{\mathcal{V}}_\epsilon^{N, \delta}$. Consider the wave equation

$$\square_m h = 0, \quad (h|_{t=0}, \partial_g h|_{t=0}) = (h^0, h^1).$$

And let

$${}^m\Gamma_\mu(h) = m^{\alpha\beta} \partial_\alpha h_{\mu\beta} - \partial_\mu(\text{tr}_m h), \quad \mu = 0, 1, \dots, n.$$

Then the solution space of the linearization of harmonic gauge conditions (3.8) and (3.11) is

$$\mathcal{T}^{N, \frac{n-1}{2}+\delta} = \{(h^0, h^1) \in \rho_0^{\frac{n-1}{2}+\delta} H_b^{N+1}(\Sigma_0) \times \rho_0^{\frac{n+1}{2}+\delta} H_b^{N+1}(\Sigma_0) : {}^m\Gamma_\mu(h)|_{t=0}, \partial_t {}^m\Gamma_\mu(h)|_t = 0\}$$

Since $\square_m {}^m\Gamma_\mu(h) = {}^m\Gamma_\mu(\square_m h)$, we have the restriction map

$${}^m\mathcal{R}_{\mathcal{F}}|_{\mathcal{T}^{N, \frac{n-1}{2}+\delta}} : \mathcal{T}^{N, \frac{n-1}{2}+\delta} \longrightarrow \widetilde{\mathcal{W}}_\infty^{N, \delta}$$

is still an isomorphism. By implicit function theorem, $\mathcal{R}_{\mathcal{F}} : \widetilde{\mathcal{V}}_\epsilon^{N, \delta} \longrightarrow \widetilde{\mathcal{W}}_{C\epsilon}^{N, \delta}$ is open if ϵ is small enough. $\square$

Proof of Theorem 2: It follows from Theorem 3 and Theorem 4 directly.

8. APPENDIX

In this section, we prove some mapping property of $P = |\Delta|^{\frac{1}{2}}$ on $\mathbb{R}^n$ for $n \geq 3$ by the same method we used in [MW], where we show the mapping properties of Radon transform and Radiation field map.

Define for $u \in C_c^\infty(\mathbb{R}^n)$

$$(8.1) \quad Pu(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i(x-y) \cdot \xi} |\xi| u(y) dy d\xi.$$

Then a similar proof as for the inverse Radon transform in [He] shows that:

$$Pu = \begin{cases} \frac{(-i)^n}{2(2\pi)^{n-1}} \mathcal{M} \partial_s^n \mathcal{R} u = \frac{1}{2(2\pi)^{n-1}} \mathcal{M} \mathcal{R} (\Delta^{\frac{n}{2}} u) & n=\text{even} \\ \frac{(-i)^n}{2(2\pi)^{n-1}} \mathcal{M} \mathcal{H} \partial_s^n \mathcal{R} u = \frac{-i}{2(2\pi)^{n-1}} \mathcal{M} \mathcal{H} \partial_s \mathcal{R} (\Delta^{\frac{n-1}{2}} u) & n=\text{odd} \end{cases}$$

where $\Delta = -\sum_{i=1}^n \partial_i^2$, $\mathcal{R}$ is the Radon transform

$$(\mathcal{R}u)(s, \omega) = \int_{x \cdot \omega = s} u(x) dH$$

with dH the induced Lebesgue measure on hyperplane $x \cdot \omega = s$ and $\mathcal{M}, \mathcal{H}$ are defined as follows: for $v \in C_c^\infty(\mathbb{R} \times \mathbb{S}^{n-1})$,

$$\begin{aligned} (\mathcal{M}v)(x) &= \int_{\mathbb{S}^{n-1}} v(x \cdot \omega, \omega) d\omega, \\ (\mathcal{H}v)(s, \omega) &= \frac{i}{\pi} \lim_{(\epsilon, N) \rightarrow (0, \infty)} \int_{\epsilon < |s-s'| < N} (s-s')^{-1} v(s', \omega) ds'. \end{aligned}$$

Proposition 8.1. *For $n \geq 3$ and $\lambda \in (0, n-1)$, P extends to an isomorphism:*

$$P : \rho_0^\lambda H_b^N(\overline{\mathbb{R}^n}) \longrightarrow \rho_0^{\lambda+1} H_b^{N-1}(\overline{\mathbb{R}^n}).$$

satisfying $P^2 = \Delta$.

Proof. By the mapping property of Δ , we only need to show the above map is continuous. For $\Re \lambda \in (0, n-1)$, by a similar proof as in [MW], P extends to a continuous map

$$\rho^\lambda C^\infty(\overline{\mathbb{R}^n}) \longrightarrow \rho^{\lambda+1} C^\infty(\overline{\mathbb{R}^n}) + \rho^{n+1} C^\infty(\overline{\mathbb{R}^n})$$

which restricts to the boundary to define

$$\begin{aligned} B(\tau)f &= \rho^{-\tau-1} P \rho^\tau \tilde{f}|_{\rho=0} : C^\infty(\mathbb{S}^{n-1}) \longrightarrow C^\infty(\mathbb{S}^{n-1}) \\ \text{where } \tilde{f} &\in C^\infty(\overline{\mathbb{R}^n}), \tilde{f}|_{\rho=0} = f. \end{aligned}$$

Here $B(\tau)$ is a meromorphic family of semiclassical pseudodifferential operator with simple poles at $(n + \mathbb{N}_0) \cup (-\mathbb{N})$. Moreover, let $A(\tau) : C^\infty(\mathbb{S}^{n-1}) \longrightarrow C^\infty(\mathbb{S}^{n-1})$ defined by the Schwartz kernel $(\theta \cdot \omega)_+^{\tau-n}$. Then

$$B(\tau) = \begin{cases} C_n(\tau) A(n-1-\tau) A(\tau) & n = \text{even} \\ \frac{i}{\pi} C_n(\tau) [-C_+(\tau+1) A(n-1-\tau) + C_-(\tau+1) \Theta A(n-1-\tau)] A(\tau) & n = \text{odd} \end{cases}$$

where Θ is the antipodal map and

$$C_n(\tau) = \frac{(-i)^n}{(2\pi)^{n-1}} \prod_{i=0}^{n-1} (\tau - i), \quad C_\pm(\tau+1) = \int_0^\infty (1 \mp t)^{-1} t^{-\tau-1} dt.$$

Hence $\chi(B(\tau))(\theta, \theta') = B_0(\theta, \theta') + B_{sc}(\theta, \theta') + B_\pm(\theta, \pm\theta')$ where

$$B_0 \in \eta^{-1} \Psi_{sc}^{-n}(\mathbb{S}^{n-1}), \quad B_{sc} \in \eta^{-1} \Psi_{sc}^1(\mathbb{S}^{n-1}), \quad B_\pm \in \mathcal{S}(\mathbb{R}_{\text{Im}\tau}, \Psi^1(\mathbb{S}^{n-1}))$$

with $\eta = \frac{1}{\text{Im}\tau}$ the semiclassical parameter as $\text{Im}\tau \rightarrow 0$. From standard estimates on the boundedness of semiclassical families,

$$B(\tau) : H^N(\mathbb{S}^{n-1}) \cap (1 + |\text{Im}\tau|)^{-1} H^{N-1}(\mathbb{S}^{n-1}) \longrightarrow H^{N-1}(\mathbb{S}^{n-1})$$

is continuous. By Mellin transform and taking $\alpha = \text{Re}\tau$, we finish the proof. $\square$

REFERENCES

- [CB1] Y. Choquet-Bruhat, *Theorem d'Existence pour certains systemes d'equations aux derivees partielles nonlin-eaires*. Acta. Math. **88**(1982), 141-225.
- [CB2] Y. Choquet-Bruhat, *The null condition and asymptotic expansions for the Einstein's equations*, Ann. Phys, (Leipzig) **9**(2000), 258-266.
- [CBG] Y. Choquet-Bruhat, R.P. Geroch, *Global aspects of the Cauchy problem in General Relativity*, CMP **14**(1969), 329-335.
- [Ch1] D. Christodoulou, *Global solutions of nonlinear hyperbolic equations for small initial data*, Comm. Pure Appl. Math. **39**(1986), 267-282.
- [CK] D. Christodoulou, S. Klainerman, *The global Nonlinear Stability of the Minkowski Space*, Princeton Mathematical Series, **41**(1993), Princeton University Press.
- [Fr] F.G. Friedlander, *Radiation fields and hyperbolic scattering theory*, Math. Proc. Cambridge Philos. Soc. **88** (1980), no.3, 483-515.
- [He] S. Helgason, *Radon Transform*, Birkhauser Boston, 1999.
- [Ho1] L. Hörmander, *Lectures on Nonlinear Hyperbolic differential equations*, Mathematiques and Applications, Vol.**26**(1997), Springer, Berlin.
- [Kl] S. Klainerman, *The null condition and global existence to nonlinear wave equations*. Lectures in Applied Mathematics **23**(1986), 293-329.
- [LR1] H. Lindblad, I. Rodnianski, *Global existence for the Einstein vacuum equations in wave coordinates*, Comm. Math. Phys. **256**(2005), No.1, 43-110.
- [LR2] H. Lindblad, I. Rodnianski, *The global stability of the Minkowski space-time in harmonic gauge*, arXiv:math/0411109.
- [Me] R. B. Melrose, *Differential analysis on manifolds with corners*, in preparation.
- [Me2] R. B. Melrose, *Transformation of boundary problems*, Acta **147**(1981), 149-236.
- [MW] R. B. Melrose, F. Wang, *Radon transform and Radiation field*, in preparation.